

N d 6 0



H



**Advil**  
children  
2-12 years  
Pain & Fever Relief

**Medicine Information**

**Active Ingredient**  
Ibuprofen 20 mg/mL

**Uses**  
Advil Children 2-12 years provides temporary relief of pain associated with:  
• Earache • Toothache • Cold and Flu • Headache  
• Minor Aches • Sore Throat • Sprains and Strains  
Advil Children 2-12 years reduces fever

**Warnings**  
**Do not use this product:**  
• At all in infants under 3 months and/or weighing less than 6 kg  
• If your child has a current or past history of a stomach ulcer or bleeding disorder  
• If your child has impaired kidney function or heart failure  
• If your child is allergic to ibuprofen, aspirin or other anti-inflammatory medicines or any other components of this product  
• If you are pregnant, or trying to become pregnant

**Ask your doctor before use if:**  
• Your child suffers from asthma, urticaria or rhinitis  
• Your child has indigestion, stomach pain, high blood pressure, heart or liver disease  
• Your child has not been taking fluids or has lost fluids due to vomiting or diarrhoea  
• Your child is taking other medicines regularly or medicines containing ibuprofen, aspirin or other anti-inflammatory ingredients or blood thinners  
• Your child is taking aspirin for heart attack or stroke because ibuprofen may decrease this benefit of aspirin  
• Your child is under doctor's care for a serious condition  
• You are breastfeeding  
• Your child is 3 months to 2 years of age

**While using this product:**  
• Stop use and ask your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days  
• Stop use and ask your doctor immediately if you see blood in the vomit or bloody or black stools  
• Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage  
• Side effects may be minimised by using for the shortest duration necessary  
• If an overdose is taken or suspected, ring the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766), discontinue use & seek professional advice immediately

**Contains:** benzoates, hydroxybenzoates, sulfites, saccharin, and sugar alcohols (18 g/60 mL). Products containing sugar alcohols may have a laxative effect or cause diarrhoea

Continued on panel to the right...>

**Advil**  
children  
2-12 years  
Pain & Fever Relief

PHARMACY MEDICINE  
KEEP OUT OF REACH OF CHILDREN

PHARMACY MEDICINE  
KEEP OUT OF REACH OF CHILDREN

HALEON

**Medicine Information (continued)**

**Directions for use based on 5-10 mg/kg/dose**

Children	Ave. body weight	How much	Time between each dose	Maximum
2 to under 4 years	12 - 16 kg	5 mL	6 - 8 hours	4 doses in 24 hours
4 to under 7 years	16 - 22 kg	7.5 mL	6 - 8 hours	4 doses in 24 hours
7 to under 10 years	22 - 32 kg	10 mL	6 - 8 hours	4 doses in 24 hours
10 to 12 years	32 - 41 kg	15 mL	6 - 8 hours	4 doses in 24 hours

• Shake bottle well before use  
• A measuring device is included for accurate dose measurement. Wash measuring device after each use

**Other information:**  
• Store below 25°C. Protect from light  
• Do not use if tamper evident seal is broken or missing

**Haleon**  
Sydney, NSW, Australia  
AU: 1800 028 533  
www.advil.net.au  
Trade Marks are owned by or licensed to the Haleon group of companies.  
© 2023 Haleon group of companies or its licensor.

Ibuprofen 20 mg/mL  
up to **8hr Fever Relief**  
strawberry banana flavour  
colour free  
sugar free  
great tasting

Dosing Device Inside  
200 mL Oral Liquid Suspension

AUST R 178433

Ibuprofen 20 mg/mL  
up to **8hr Fever Relief**  
strawberry banana flavour  
colour free  
sugar free  
great tasting

Dosing Device Inside  
200 mL Oral Liquid Suspension

AUST R 178433

C200-APFC-6



9 310488 000265



C9PN