





# Medicine Information (continued)

- White using this product:

  Stop use and ask your doctor immediately if you see an altergic reaction including skin reddening, rash of bilsters or if pain or fever gets worse or lasts for more than a few days. Stop use and ask your doctor immediately if you see blood in the youn't or bloody or hack stools. Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart dissease, stroke or liver damage.

  Side effects may be minimised by using for the shortest duration necessary. If an overdose is taken or suspected, ring the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766), discontinue use & seek professional advice immediately

  Contains: benzoates, hydroxybenzoates, suffites, sugars

# Directions for Use:

# Dosage based on 5 - 10 mg/kg/dose

	Age			Time Between Each Dose	Maximum
	3 to under 6 months	6 to 8 kg	1.5 mL	6 <b>-</b> 8 hours	4 doses in 24 hours
	6 to under 12 months	8 to 10 kg	2 mL	6 <b>-</b> 8 hours	4 doses in 24 hours
	1 to 2 years	10 to 12 kg	2.5 mL	6 <b>-</b> 8 hours	4 doses in 24 hours

- Shake bottle well before use
   A measuring device is included for accurate dose measurement Wash measuring device after each use
   For children 3 months to 12 months give only on medical advice

- Store below 25°C
   Do not use if tamper evident seal is broken or missing

Haleon Sydney, NSW, Australia AU: 1800 028 533

# @ www.advil.net.au

Trade Marks are owned by or licensed to the Haleon group of companies.

© 2023 Haleon group of companies or its licensor.

# PHARMACY MEDICINE

# infant

3 months - 2 years

# Pain & Fever Relief

Ibuprofen 40 mg/mL #8hr Fever Relief



infant



3 months - 2 years Pain & Fever Relief

Ibuprofen 40 mg/mL #8hr Fever Relief

grape flavour

Dosing
Device Inside
40 mL
Oral Liquid
Suspension

Concentrated Formula

# Medicine Information

# **Active Ingredient** Ibuprofen 40 mg/mL

HAL≣ON

- Uses
  Advil Infant 3 months 2 years provides temporary relief of pain associated with:

   Earache Tenthing Cold and Flu
   Headache Minor Aches Sprains and Strains
   Immunisation
  Advil Infant 3 months 2 years reduces fever.

- Warnings
  Do not use this product:

  In infants under 3 months and/or weighing less than 6 kg
  If your Child has a current or past history of a stomach
  ulce or o bleeding disorder
  If your child has impaired kidney function or heart failure
  If your child is allergist to bupprofen, aspirin or other
  anti-inflammatory medicines or any other components
  of this product
  If you are pregnant, or trying to become pregnant

- Nyour de pregnant, or vying to become pregnant.

  Ask your doctor before use if:
   Your child suffers from asthma, urticaria or rhinitis
   four child suffers from asthma, urticaria or rhinitis
   four child has indigestion, stomach pain, high blood pressure, heart or liver disease
   four child has not been taking fluids or has lost fluids due to vomiting or diarrhoea
   four child is sand the medicines regularly or medicines containing bupprofen, aspirin or other anti-inflammatory ingredients or blood thinners
   four child is taking aspirin for heart attack or stoke because bupprofen may decrease this benefit
   of aspirin
   four child is under doctor's care for a serious condition
   four are breastfeeding
- - Continued on panel to the right...▶

C40-APFI-6

