

# New Formulation



**Panadol Children 1-5 years**  
ORANGE Flavour  
Colour free & sugar free suspension

**PHARMACY MEDICINE**  
KEEP OUT OF REACH OF CHILDREN  
HALEON

**Panadol Children 1-5 years**  
Paracetamol 24 mg/mL  
Fast pain relief

**NEW FORMULA & IMPROVED ORANGE FLAVOUR**  
100 mL Colour free & sugar free suspension

**Active ingredients**  
This suspension contains PARACETAMOL 24 mg/mL.

**What this medicine is used for**  
Effective temporary relief of fever and pain associated with:  
• Teething • Irritation • Earache  
• Headache • Cold flu symptoms

**Warnings**  
**Do not use this medicine**  
• If your child is allergic to paracetamol  
• If your child is using other medicines containing paracetamol, except on medical advice  
• For more than 48 hours at a time, except on medical advice.  
• If any of the seals on this packaging are broken  
• If the package use-by date has expired  
**Ask a doctor before use if your child**  
• Has liver or kidney problems  
• Weighs less than 10 kg or more than 41 kg  
**Stop use and see your doctor if your child**  
Has an allergic skin reaction, shortness of breath or wheezing after taking this medicine.

**While using this product**  
If an overdose is seen or suspected, ring the Poisons Information Centre (AUST: 13 11 26) or go to the hospital immediately, even if your child feels well because of the risk of delayed serious liver damage if left untreated. Keep to the recommended dose.  
**Contents:** hydroxybenzoin, supralin, metformin (27 g/104 mL) and sorbitol (14 g/104 mL). Products containing metformin and sorbitol may have a laxative effect or cause diarrhoea.

**Directions for use**  
• Read and retain the carton  
• Shake the bottle well

**Directions for use (continued)**

- Follow instructions for using the provided dosing device to accurately measure.
- Measure the correct dose based on the dosage table.
- Calculate correct dose based on child's weight.
- If weight is unknown, use the child's age.
- Repeat 4-6 hourly if required (maximum 4 times within 24 hours).
- Can be given with water or fruit juice if preferred by your child.
- Further information is available via the QR code on the pack.

Age	Weight	Dose
1-2 Yrs	10-12 kg	6-8 mL
2-3 Yrs	12-14 kg	8-10 mL
3-4 Yrs	14-16 kg	9-10 mL
4-5 Yrs	16-18 kg	10-11 mL
5-6 Yrs	18-20 kg	11-13 mL
6-7 Yrs	20-22 kg	12-14 mL
7-8 Yrs	22-25 kg	14-16 mL
8-9 Yrs	24-28 kg	16-18 mL
9-10 Yrs	26-32 kg	18-20 mL
10-11 Yrs	30-36 kg	20-23 mL
11-12 Yrs	36-41 kg	23-26 mL

**After using this product**  
• Wash dosing device  
• Wipe the neck of the bottle clean  
• Push the cap down and turn it to close cap tightly, then turn backwards until you hear clicking sounds.

Store below 30°C. Do not refrigerate.

**Haleon**  
NSW, Australia  
Trade Marks are owned by or licensed to the Haleon group of companies.  
©2025 Haleon group of companies or its licensor.

**AU: 1800 028 533**  
www.chil@panadol.com.au  
Manufactured by OUC (Thailand) Limited, Thailand

**Using the dosing device**

- Push the plunger fully up to close the cap. Push the cap down and turn it to close cap tightly, then turn backwards until you hear clicking sounds.
- Turn the bottle upside down and gently pull the correct dose for your child. The correct dose is at the point where the widest sides of the plunger meet the correct ml mark on the barrel of the dosing device (ie. where the liquid ends as shown by ②).
- Turn the bottle upright and remove the device from the neck of the bottle by gently twisting it.

**Effective pain and fever relief associated with:**  
• Headache • Earache • Irritation • Teething  
• Cold flu symptoms • Teething

• Colour free  
• Sugar free

**Directions for use (continued)**

• Read and retain the carton  
• Shake the bottle well

991017600



9 500673 896063

1 Ver A

Old  
Formulation

0 8 1 1

**1-5 YEARS**  
Orange Flavour  
Colourfree

**ACTIVE INGREDIENTS**

This suspension contains PARACETAMOL 24 mg/mL

**WHAT THIS MEDICINE IS USED FOR**

Effective temporary relief of fever and pain associated with:

- Teething • Immunisation • Earache
- Headache • Cold & flu symptoms

**WARNINGS**

**Do not use this medicine**

- If your child is using other medicines containing paracetamol, except on medical advice
- For more than 48 hours at a time, except on medical advice. Keep to the recommended dose.
- If any of the seals on this packaging are broken
- If the package use-by date has expired

**Ask a doctor before use if your child**

- Has fever or kidney problems
- Weighs less than 10 kg or more than 41 kg

**Stop use and see your doctor if your child**

Has an allergic skin reaction, shortness of breath or wheezing after taking this medicine.

**While using this product**

If an overdose is seen or suspected, ring the Poisons Information Centre (AUST: 13 11 26) or go to the hospital immediately, even if your child feels well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose.

Contains hydrobenzoinone, sorbitol, sorbitol (1.96 g/104 mL) and maltitol (38.59 g/104 mL) which may have a laxative effect or cause diarrhoea.

**DIRECTIONS FOR USE**

- Please read and retain the carton
- Shake the bottle well
- Use the dosing device provided to accurately measure
- Measure the correct dose based on the dosage table
- Calculate correct dose based on child's weight, if weight is unknown, use the child's age
- Repeat 4–5 hourly if required (maximum 4 times within 24 hours)
- Can be given with water or fruit juice if preferred by your child

Further information is available via the QR code on the pack.

Continued →

**DIRECTIONS FOR USE (continued)**

Age	Average Weight	Dose
1–2 Yrs	10–12 kg	6–8 mL
2–3 Yrs	12–14 kg	8–9 mL
3–4 Yrs	14–16 kg	9–10 mL
4–5 Yrs	16–18 kg	10–11 mL
5–6 Yrs	18–20 kg	11–13 mL
6–7 Yrs	20–22 kg	13–14 mL
7–8 Yrs	22–25 kg	14–16 mL
8–9 Yrs	25–28 kg	16–18 mL
9–10 Yrs	28–32 kg	18–20 mL
10–11 Yrs	32–36 kg	20–23 mL
11–12 Yrs	36–41 kg	23–26 mL

**Using the dosing device**

- 1 Push the plunger fully in to close the dosing device, then insert it firmly into the neck of the bottle.
- 2 Turn the bottle upside down and gently pull the correct dose for your child. The correct dose is at the point where the widest sides of the plunger meets the correct mL mark on the barrel of the dosing device (ie, where the liquid ends as shown by ①).
- 3 Turn the bottle upright and remove the device from the neck of the bottle by gently twisting it.



**After using the medicine**

- Hand wash dosing device
- Wipe the neck of the bottle clean
- Push the cap down and turn it to close cap tightly, then turn backwards until you hear clicking sounds.

Store below 30°C. Do not refrigerate.

This medicine contains: No artificial colours

Haleon  
Sydney, NSW, Australia  
Trade Marks are owned by or licensed to the Haleon group of companies.  
©2022 Haleon group of companies or its licensor.  
AU: 1800 028 533  
www.childrenspanadol.com.au

**1-5 YEARS**  
Orange Flavour  
Colourfree

**PHARMACY MEDICINE**  
KEEP OUT OF REACH OF CHILDREN

HALEON



**1-5 YEARS**  
Paracetamol 24 mg/mL

**FAST PAIN RELIEF**  
Reduces fever  
Gentle on stomach



**1-5 YEARS**  
Paracetamol 24 mg/mL

**FAST PAIN RELIEF**  
Reduces fever  
Gentle on stomach

Orange Flavour  
Colourfree suspension 100 mL

**H**

9 300673 916792  
7000644

1 1 8 0