

0 8 1 1

1-5 YEARS
Orange Flavour
Colourfree

ACTIVE INGREDIENTS

This suspension contains PARACETAMOL 24 mg/mL

WHAT THIS MEDICINE IS USED FOR

Effective temporary relief of fever and pain associated with:
• Teething • Immunisation • Earache
• Headache • Cold & flu symptoms

WARNINGS

Do not use this medicine

- If your child is using other medicines containing paracetamol, except on medical advice
- For more than 48 hours at a time, except on medical advice. Keep to the recommended dose.

- If any of the seals on this packaging are broken
- If the package use-by date has expired
- Ask a doctor before use if your child:
 - Has liver or kidney problems
 - Weighs less than 10 kg or more than 41 kg

Stop use and see your doctor if your child

Has an allergic skin reaction, shortness of breath or wheezing after taking this medicine

While using this product

If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 13 11 26) or go to the hospital immediately, even if your child feels well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose.

Contains: hydroxybenzoates, sucralose, sorbitol (11.96 g/104 mL) and maltitol (38.58 g/104 mL) which may have a laxative effect or cause diarrhoea.

DIRECTIONS FOR USE

- Please read and retain the carton
- Shake the bottle well
- Use the dosing device provided to accurately measure
- Measure the correct dose based on the dosage table
- Calculate correct dose based on child's weight. If weight is unknown, use the child's age.
- Repeat 4-6 hourly if required (maximum 4 times within 24 hours)
- Can be given with water or fruit juice if preferred by your child
- Further information is available via the QR code on the pack.

Continued →

DIRECTIONS FOR USE (continued)

Age	Average Weight	Dose
1-2 Yrs	10-12 kg	6-8 mL
2-3 Yrs	12-14 kg	8-9 mL
3-4 Yrs	14-16 kg	9-11 mL
4-5 Yrs	16-18 kg	10-11 mL
5-6 Yrs	18-20 kg	11-13 mL
6-7 Yrs	20-22 kg	13-14 mL
7-8 Yrs	22-25 kg	14-16 mL
8-9 Yrs	25-28 kg	16-18 mL
9-10 Yrs	28-32 kg	18-20 mL
10-11 Yrs	32-36 kg	20-23 mL
11-12 Yrs	36-41 kg	23-26 mL

Using the dosing device

- 1 Push the plunger fully in to close the dosing device; then insert it firmly into the neck of the bottle.
- 2 Turn the bottle upside down and gently pull the correct dose for your child. The correct dose is at the point where the widest sides of the plunger meets the correct mL mark on the barrel of the dosing device (i.e. where the liquid ends as shown by ⊕).
- 3 Turn the bottle upright and remove the device from the neck of the bottle by gently twisting it.



After using this medicine

- Hand wash dosing device
- Wipe the neck of the bottle clean
- Push the cap down and turn it to close cap tightly, then turn backwards until you hear clicking sounds

Store below 30°C. Do not refrigerate.

This medicine contains: No artificial colours

Haleon
Sydney, NSW, Australia
Trade Marks are owned by or licensed to the Haleon group of companies.

©2023 Haleon group of companies or its licensor.

AU: 1800 028 533

www.childrenspanadol.com.au



9 300673 916792

9000644



1 1 8 0

1-5 YEARS
Orange Flavour
Colourfree

Panadol Children
1-5 YEARS
Orange Flavour
Colourfree

e-leaflet
Please scan this code for more information about this medicine.

PHARMACY MEDICINE
KEEP OUT OF REACH OF CHILDREN
HALEON

Panadol Children

1-5 YEARS
Paracetamol 24 mg/mL
FAST PAIN RELIEF
Reduces fever
Gentle on stomach

Orange Flavour
Colourfree suspension
100 mL

PHARMACY MEDICINE
KEEP OUT OF REACH OF CHILDREN
HALEON

Panadol Children

1-5 YEARS
Paracetamol 24 mg/mL
FAST PAIN RELIEF
Reduces fever
Gentle on stomach

Orange Flavour
Colourfree suspension
100 mL