



Panadol Children 5-12 years



•-tablet
Please scan this code for more information about this medicine.

Tutti Frutti flavour
Colour free & sugar free suspension

Panadol Children 5-12 years

Tutti Frutti flavour



Gentle on stomach



•Colour free
•Sugar free



Effective pain and fever relief associated with:
• Headache • Earache • Immunisation
• Cold & flu symptoms • Teething

Using the dosing device

- 1 Push the plunger fully in to close the dosing device, then insert it firmly into the neck of the bottle.
- 2 Turn the bottle upside down and gently pull the correct dose for your child. The correct dose is at the point where the red sides of the plunger meets the correct mL mark on the barrel of the dosing device (i.e. where the liquid ends as shown by 0.5).
- 3 Turn the bottle upright and remove the device from the neck of the bottle by gently twisting it.



PHARMACY MEDICINE
KEEP OUT OF REACH OF CHILDREN
HALEON

Panadol Children

5-12 years

Paracetamol 48 mg/mL

Fast pain relief

- Reduces fever
- Gentle on stomach



Tutti Frutti flavour
Colour free & sugar free suspension

200 mL

Active ingredients

This suspension contains PARACETAMOL 48 mg/mL.

What this medicine is used for

Effective temporary relief of fever and pain associated with:
• Teething • Immunisation • Earache
• Headache • Cold & flu symptoms

Warnings

- Do not use this medicine**
- If your child is allergic to paracetamol
 - If your child is using other medicines containing paracetamol, except on medical advice
 - For more than 48 hours at a time, except on medical advice.
 - Keep to recommended dose
 - If any of the seals on this packaging are broken
 - If the package use-by date has expired
 - Ask a doctor before use if your child
 - Has liver or kidney problems
 - Has liver or kidney problems
 - Weighs less than 10 kg or more than 41 kg
 - Stop use and see your doctor if your child has an allergic skin reaction, shortness of breath or wheezing after taking this medicine
- While using this product**
- If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 1311 26) or go to the hospital immediately, even if your child feels well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose.
- Contains:** hydroxybenzoates, sucralose, maltitol (0.5 g/52 mL) and sorbitol (7 g/52 mL). Products containing maltitol and sorbitol may have a laxative effect or cause diarrhoea.

Directions for use

- Read and retain the carton
- Shake the bottle well

Directions for use (continued)

- Follow instructions for using the provided dosing device to accurately measure
- Measure the correct dose based on the dosage table
- Calculate correct dose based on child's weight. If weight is unknown, use the child's age.
- Repeat 4-6 hourly if required (maximum 4 times within 24 hours)
- Can be given with water or fruit juice if preferred by your child
- Further information is available via the QR code on the pack.

Age	Average Weight	Dose
1-2 Yrs	10-12 kg	3-4 mL
2-3 Yrs	12-14 kg	4 mL
3-4 Yrs	14-16 kg	4-5 mL
4-5 Yrs	16-18 kg	5 mL
5-6 Yrs	18-20 kg	6 mL
6-7 Yrs	20-22 kg	6-7 mL
7-8 Yrs	22-25 kg	7-8 mL
8-9 Yrs	25-28 kg	8-9 mL
9-10 Yrs	28-32 kg	9-10 mL
10-11 Yrs	32-36 kg	10-11 mL
11-12 Yrs	36-41 kg	11-13 mL

After using this product

- Hand wash dosing device
- Wipe the neck of the bottle clean
- Push the cap down and turn it to close cap tightly, then turn backwards until you hear clicking sounds

Store below 30°C. Do not refrigerate.

Haleon
Sydney, NSW, Australia
Trade Marks are owned by or licensed to the Haleon group of companies.
©2025 Haleon group of companies or its licensor.
AUS: 1800 026 533
www.childrenspanadol.com.au
Manufactured by OUC (Thailand) Limited, Thailand



99107665



9 300673 897084

1 Ver. 4