



Panadol Children
Suppositories
6 Months to 5 Years

ACTIVE INGREDIENT
Each suppository contains Paracetamol 125 mg

INDICATIONS
For the relief of fever, pain, cold, influenza, cough, sore throat, earache, teething, toothache, headache, and vomiting for effective temporary relief of fever and pain associated with:

- Teething
- Immunisation
- Earache
- Headache
- Cold & flu symptoms

WARNINGS

Do not use

- If your child is allergic to paracetamol
- If your child is using other medicines containing paracetamol
- For the first time in children aged 6 months to 5 years
- If the child has been taking paracetamol for more than 48 hours at a time, except on medical advice
- If the package seals are broken or the use-by-date has expired
- Until you have read the enclosed leaflet carefully

Ask a doctor before use if your child

- Weighs less than 9 kg or more than 18 kg
- Has an allergic skin reaction, shortness of breath or wheezing after using this medicine

Stop use and see a doctor if your child

- Has an allergic skin reaction, shortness of breath or wheezing after using this medicine

While using this medicine

- If an overdose is taken or suspected, ring the Poisons Information Centre (AUST) 13 11 25 (N.Z.) 0800 764 736 (N.Z.) immediately. Do not induce vomiting because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose. Keep out of sight and reach of children.

DIRECTIONS FOR USE

Paracetamol is only to be swallowed. Do not chew, suck, or suck on the suppository. For ease of insertion, the suppository can be moistened just before insertion. Insert into the rectum according to the dosage table below. Give the correct dose based on the child's weight. If weight is unknown, use the child's age. Weight & Dose

| | | |
|------------------|----------|------------------------|
| Age | Weight | Dose |
| 6 months-4 years | 8-16 kg | 1 (max. 4 in 24 hours) |
| 4-5 years | 16-18 kg | 2 (max. 8 in 24 hours) |

Further information is available via the QR code on the pack.



Panadol Children
Suppositories
6 Months to 5 Years

10 suppositories

PHARMACY MEDICINE
KEEP OUT OF REACH OF CHILDREN
HALEON

Panadol Children

Suppositories
6 Months to 5 Years

Paracetamol 125 mg

EFFECTIVE PAIN RELIEF
10 suppositories

AUST R 3327B

Store below 25°C.
Haleon
Sydney, NSW, Australia

Trade Marks are owned by or licensed to the Haleon group of companies.
© 2023 Haleon group of companies or its licensors.

AU: 1800 028 533
www.childrenspanadol.com.au

e-leaflet
Please scan this code for more information about this medicine.

CARTON RECYCLE



5059896 AU/NZ

EXP:
BATCH:



HALEON

Panadol Children Suppositories

6 Months - 5 Years & 5 - 12 Years

125 mg paracetamol & 250 mg paracetamol

How to use it

If possible, ask your child to go to the toilet and empty their bowels before using the suppository.

Suppositories work best if the bowels are emptied.

Tear the blister along the perforated line between each suppository.

You can then remove the suppository by pulling each side of the blister apart from the top.

Follow these steps to use a suppository:

1. Wash your hands thoroughly with soap and water.
2. Feel the suppository while it is still in the foil.
3. If it feels soft, keep it in the foil, and chill it in the fridge or by holding it under cold water for a few minutes. Do not remove the foil wrapper while you are chilling it.
4. Put on a disposable glove, if desired. They are available from pharmacies.
5. Remove all of the foil wrapper from the suppository.
6. Moisten the suppository by dipping it briefly in cool water.
7. Lie your child on their side and raise their knee to their chest.
8. Push the suppository gently, blunt end first, into their rectum, or back passage.
9. Keep your child lying down for a few minutes so that the suppository dissolves.
10. Throw away used materials and wash your hands thoroughly.

Trade Marks are owned by or licensed to the Haleon group of companies.
©2026 Haleon group of companies or its licensor.