Each Panadol Rapid caplet CONTAINS PARACETAMOL 500 mg



USE ONLY IF SEAL IS INTACT



Panadol Rapid

ACTIVE INGREDIENTS

Each caplet CONTAINS PARACETAMOL 500 mg

WHAT THIS MEDICINE IS USED FOR

Fast, effective temporary relief of pain and discomfort associated with: • Headache/Tension headache • Migraine headache • Muscular aches • Period pain • Toothache • Cold & flu symptoms. **Reduces fever**

WARNINGS

Do not use Panadol • If you are allergic to paracetamol • If using other medicines containing paracetamol • For children below age 12 • For more than 48 hours at a time for children aged 12–17 except on medical advice • For more than a few days at a time in adults except on medical advice • If any of the seals on this packaging are broken • If the package use-by date has expired • Until you read the enclosed leaflet carefully

Ask a doctor before use • If you have liver or kidney problems • If you are taking warfarin (a medicine used to thin the blood).

Stop use and see your doctor immediately • If you have an allergic skin reaction, shortness of breath or wheezing after taking Panadol.

While using this product • If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 13 11 26; NZ: 0800 764 766) or go to the hospital immediately, even if you feel well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose. Two caplets contain 352 mg (15.3 mmol) of sodium. This should be taken into account by people on a low sodium diet. Keep out of reach & sight of children.

DIRECTIONS FOR USE

Age:	caplets	How often
12–Adult	2	every 4–6 hrs with water as required (maximum 8 caplets in 24 hrs)

DIRECTIONS FOR USE

Store below 30°C.

Suitable for: People with stomach ulcers • Breastfeeding mothers

Haleon
Sydney, NSW, Australia & Auckland, New Zealand

9 300673 898449

AU: 1800 028 533
NZ: 0800 540 144
www.panadol.com.au
www.panadol.co.nz

Trade Marks owned or licensed by Haleon.
©2024 Haleon or licensor.

302153

CARTON RECYCLE

Dispose of blister section in the bin.

Panadol Rapid

8 caplets

AUST R 332528

HALEON

Panadol

Rapid

Paracetamol 500 mg

Fast Pain Relief

8 caplets capsule shaped tablets

PANADOL Rapid Caplets
500 mg Paracetamol

1. WHAT PANADOL RAPID IS AND WHAT IS IT USED FOR

PANADOL Rapid caplets are used for the fast effective temporary relief of pain and discomfort associated with: headache, muscular aches, period pain, toothache, migraine headache, colds & flu and tension headache.
The active ingredient is paracetamol which works to stop the pain messages from getting through to the brain and also reduces fever. Always use the lowest effective dose to relieve symptoms for the shortest period of time necessary.

2. CHECK BEFORE YOU TAKE PANADOL RAPID

Do not take PANADOL Rapid caplets:
• If you have had an allergic reaction to paracetamol or any of the ingredients in this product.
• If you are taking other medicines containing paracetamol.
• If you are under 12 years.

Ask your doctor before you take this medicine if you:
• Have **liver or kidney** problems, including alcoholic liver disease, are **underweight** or **malnourished** or **regularly drink alcohol.**
• You have a severe infection as this may increase the risk of metabolic acidosis. Signs of metabolic acidosis include: deep, rapid, difficult breathing; feeling sick (nausea) or being sick (vomiting); loss of appetite.
Please see your doctor if your symptoms do not improve. Keep out of sight and reach of children.

If you are taking other medicines
Talk to your doctor or pharmacist before taking these caplets if you are taking other medicines, particularly **metoclopramide** (for **nausea** [feeling sick] or **vomiting** [being sick]; **cholestyramine** (to **lower cholesterol**). If you **take blood thinning medicines** (**anticoagulants** e.g. **warfarin**) and you need to take a pain reliever regularly, **talk to your doctor** because of the risk of bleeding.

Pregnancy & Breastfeeding
Talk to your doctor before taking PANADOL Rapid caplets if you are pregnant. You can take this product while breastfeeding.

Important information about some of the ingredients
Two PANADOL Rapid caplets contain 352 mg sodium. If you are on a low sodium diet, ask your doctor about using this medicine before you take it.

3. SIDE EFFECTS

Like all medicines, PANADOL Rapid caplets can have side effects, but not everybody gets them. Stop taking this medicine and tell your doctor immediately if you experience:
• Shortness of breath
• Wheezing or difficulty breathing
• Swelling of the face, lips, tongue, throat or other parts of the body
• Rash, peeling, itching or hives on the skin or mouth ulcers
• Unexplained bruising or bleeding
These side effects are very rare for low doses of this medicine and when used for a short period of time.
If any of the previous side effects happen, tell your pharmacist or doctor immediately or go to Accident and Emergency at your nearest hospital.
Tell your pharmacist or doctor if you notice anything that is making you feel unwell.
Other side effects not listed above may also occur in some people.

4. FURTHER INFORMATION

PANADOL Rapid caplets contain 500 mg of paracetamol as the active ingredient.
They also contain: sodium bicarbonate, cellulose - microcrystalline, starch - pregelatinised maize, sodium starch glycolate, starch - maize, water - purified, magnesium stearate, hypromellose, colloidal anhydrous silica, povidone, titanium dioxide, polydextrose, glycerol triacetate, carnauba wax and macrogol 8000.
PANADOL Rapid caplets are a white to off-white capsule shaped film coated tablets with flat edges. Tablets debossed with "P" on one side and "---" on the other side. They come in blister packs of 16, 32 & 48 caplets and a 'Handipak' of 8 caplets.

Haleon
Sydney, NSW, Australia & Auckland, New Zealand
AUST R 332528 (PANADOL Rapid caplets)
Date of preparation: November 2025.