Superseded Artwork

17606

Panadol

Rapid Soluble

16 effervescent tablets

HALEON

Panadol

Rapid Soluble

Paracetamol 500 mg

Fast Pain Relief

AUST R 15509

16 effervescent tablets

LEMON FLAVOUR

Panadol

Rapid Soluble

PNDLCTN16

16 effervescent tablets

Rapid Soluble

Panadol

## ACTIVE INGREDIENTS

Each effervescent tablet CONTAINS PARACETAMOL 500 mg

## WHAT THIS MEDICINE IS USED FOR

Fast effective temporary relief of pain and discomfort associated with:

- Headache/Tension headache • Migraine headache
- Muscular aches • Period pain • Toothache
- Cold & flu symptoms • Sore throat. Reduces fever

## WARNINGS

**Do not use Panadol**

- If you are allergic to paracetamol
- If using other medicines containing paracetamol
- For children below age 12 except on medical advice
- For more than 48 hours at a time for children aged 12–17 except on medical advice
- For more than a few days at a time in adults except on medical advice
- If you are PHENYLKETONURIC
- If any of the seals on this packaging are broken
- If the package use-by date has expired

**Ask a doctor before use**

- If you have liver or kidney problems
- If you are taking warfarin (a medicine used to thin the blood)

**Stop use and see your doctor immediately if**

You have an allergic skin reaction, shortness of breath or wheezing after taking Panadol.

**While using this product**

If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 13 11 26; NZ: 0800 764 766) or go to the hospital immediately, even if you feel well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose.

Each tablet contains, phenylalanine, saccharin sodium, sucrose and aspartame. Keep out of reach & sight of children.

Each tablet contains 425.5 mg (18.5 mmol) of sodium. This should be taken into account by those on a low sodium diet.

## DIRECTIONS FOR USE

| Age | Tablets | How often |
|---|---|---|
| 12–Adult | 2 | dissolved in a glass of water at room temperature every 4–6 hrs as required (maximum 8 tablets in 24 hrs) |

Further information is available via the QR code on the pack.

**Suitable for:**

- People with stomach ulcers • Breastfeeding mothers

9 300673 891303

Store below 30°C.

**Haleon**

Sydney, NSW, Australia & Auckland, New Zealand

**AU: 1800 028 533 NZ: 0800 540 144**

**www.panadol.com.au www.panadol.co.nz**

Trade Marks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies or its licensor.

**e-leaflet**

Please scan this code for more information about this medicine.

CARTON RECYCLE

17606

New Artwork

99101769O

Panadol

Rapid Soluble

16 effervescent tablets

HALEON

Panadol

Rapid Soluble

Paracetamol 500 mg

Fast Pain Relief

AUST R 15509

16 effervescent tablets

LEMON FLAVOUR

16 effervescent tablets

Rapid Soluble

Panadol

**ACTIVE INGREDIENTS**

Each effervescent tablet CONTAINS PARACETAMOL 500 mg

**WHAT THIS MEDICINE IS USED FOR**

Fast effective temporary relief of pain and discomfort associated with:

- Headache/Tension headache • Migraine headache
- Muscular aches • Period pain • Toothache
- Cold & flu symptoms • Sore throat. Reduces fever

**WARNINGS**

**Do not use Panadol**

- If you are allergic to paracetamol
- If using other medicines containing paracetamol
- For children below age 12 except on medical advice
- For more than 48 hours at a time for children aged 12–17 except on medical advice
- For more than a few days at a time in adults except on medical advice
- If you are PHENYLKETONURIC
- If any of the seals on this packaging are broken
- If the package use-by date has expired

**Ask a doctor before use**

- If you have liver or kidney problems
- If you are taking warfarin (a medicine used to thin the blood)

**Stop use and see your doctor immediately if**

You have an allergic skin reaction, shortness of breath or wheezing after taking Panadol.

**While using this product**

If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 13 11 26; NZ: 0800 764 766) or go to the hospital immediately, even if you feel well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose.

Each tablet contains sulfites, phenylalanine, saccharin sodium, sucrose and aspartame. Keep out of reach & sight of children.

Each tablet contains 425.5 mg (18.5 mmol) of sodium. This should be taken into account by those on a low sodium diet.

**DIRECTIONS FOR USE**

| Age | Tablets | How often |
|---|---|---|
| 12–Adult | 2 | dissolved in a glass of water at room temperature every 4–6 hrs as required (maximum 8 tablets in 24 hrs) |

Further information is available via the QR code on the pack.

**Suitable for:**

- People with stomach ulcers • Breastfeeding mothers

9 300673 891303 >

(8) EXP

Panadol

Rapid Soluble

Store below 30°C

Manufactured by : OLIC (Thailand) Limited, Thailand

**Haleon**

Sydney, NSW, Australia & Auckland, New Zealand

**AU: 1800 028 533 NZ: 0800 540 144**

**www.panadol.com.au www.panadol.co.nz**

Trade Marks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies or its licensor.

**e-leaflet**

Please scan this code for more information about this medicine.

CARTON RECYCLE

H