



**ACTIVE INGREDIENT**

Each suppository contains Paracetamol 500 mg

**WHAT THIS MEDICINE IS USED FOR**

Effective temporary relief of pain. Reduces fever.

**WARNINGS**

**Do not use Panadol Suppositories**

• If you are allergic to paracetamol • If using other medicines containing paracetamol • For children below the age of 10, except on medical advice • For more than 48 hours for children aged 10-17, except on medical advice • For more than a few days at a time in adults, except on medical advice • If any of the seals on this packaging are broken • If the package use-by date has expired • Until you read the enclosed leaflet carefully

**Ask a doctor before use**

• If you have liver or kidney problems • If you are taking warfarin (a medicine used to thin the blood)

**Stop use and tell your doctor immediately**

• If you have an allergic skin reaction, shortness of breath or wheezing after taking this medication

**While using this product**

If an overdose is taken or suspected, ring the Poisons Information Centre (AUST: 13 11 25) or go to the hospital immediately even if you feel well because of the risk of delayed, serious liver damage if left untreated. Keep to the recommended dose. Keep out of sight and reach of children.

**DIRECTIONS FOR USE**

For rectal use only - not to be swallowed.

• Tear suppository from strip, remove blister and insert suppository • For ease of insertion, the suppository can be moistened just before insertion

• Insert into the rectum according to the dosage table below every 4-6 hours as required

Age	Suppositories	How often
12 Yrs - Adult	1-2	Up to 4 times in 24 hrs as required (max 8 in 24 hrs)
10-12 Yrs	1	Up to 4 times in 24 hrs as required (max 4 in 24 hrs)

Further information is available via the QR Code on the pack.



NPAD0945

EXP:  
BATCH:



**Suppositories**

**10 suppositories**

**PHARMACY MEDICINE**  
KEEP OUT OF REACH OF CHILDREN  
HALEON



**Suppositories**

Paracetamol 500 mg

Effective temporary pain relief

**10 suppositories**

AUST R 332712

Store below 25°C.

Haleon  
Sydney, NSW, Australia &  
Auckland, New Zealand

Trade Marks are owned by or licensed to the Haleon group of companies.  
©2025 Haleon group of companies or its licensor.

AU: 1800 028 533 NZ: 0800 540 144

www.panadol.com.au www.panadol.co.nz



e-leaflet

Please scan this code for more information about this medicine.



# PANADOL Suppositories

500 mg Paracetamol

## How to use it

**If possible, go to the toilet and empty your bowels before using your suppository.**

Suppositories work best if your bowels are emptied.

Tear the blister along the perforated line between each suppository. You can then remove the suppository by pulling each side of the blister apart from the top.

## Follow these steps to use a suppository.

- 1 Wash your hands thoroughly with soap and water.
- 2 Feel the suppository while it is still in the foil.
- 3 If it feels soft, keep it in the foil, and chill it in the fridge or by holding it under cold water for a few minutes. Do not remove the foil wrapper while you are chilling it.
- 4 Put on a disposable glove, if desired. They are available from pharmacies.
- 5 Remove all of the foil wrapper from the suppository.
- 6 Moisten the suppository by dipping it briefly in cool water.
- 7 Lie on your side and raise your knee to your chest.
- 8 Push the suppository gently, blunt end first, into your rectum, or back passage.
- 9 Remain lying down for a few minutes so that the suppository dissolves.
- 10 Throw away used materials and wash your hands thoroughly.

Trade Marks are owned by or licensed to the HALEON group of companies.

©2025 HALEON group of companies or its licensor.

108



108