



1 0 9 3 7

1 0 9 3 7



Voltaren

Rapid Liquid Capsule

Diclofenac potassium 12.5 mg

What is in this leaflet

This leaflet answers some common questions about Voltaren Rapid liquid capsule. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor. All medicines have risks and benefits. Your pharmacist or doctor will be able to advise you about the risks and benefits of taking Voltaren Rapid liquid capsule.

Keep this leaflet, you may need to read it again.

Ask a doctor or a pharmacist if you need more information or advice.

What Voltaren Rapid liquid capsule is and what it is used for

Voltaren Rapid liquid capsule provides fast temporary relief from:

- muscular pain, joint pain, backache, headache, dental pain, menstrual cramps (period pain)
- painful symptoms of colds and flu (including aches and pains, sore throat pain).

Voltaren Rapid liquid capsule also reduces fever.

Voltaren Rapid liquid capsule belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs). Like all NSAIDs, Voltaren Rapid liquid capsule can relieve the symptoms of pain and inflammation; it will not cure your condition.

Voltaren Rapid liquid capsule is not addictive.

Before you take it

When you must not take it

Do not take Voltaren Rapid liquid capsule if you are allergic (hypersensitive) to:

- diclofenac, the active ingredient in Voltaren Rapid liquid capsule, or any of the ingredients listed at the end of this leaflet
- other medicines containing diclofenac
- aspirin
- any other anti-inflammatory medicines.

Many medicines used to treat headache, period pain and other aches and pains contain aspirin or anti-inflammatory medicines. If you are not sure if you are taking any of these medicines, ask your pharmacist.

Symptoms of an allergic reaction to these medicines may include:

- asthma, wheezing or shortness of breath
- swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing
- hives, itching or skin rash
- fainting.

If you are allergic to aspirin or anti-inflammatory medicines and take Voltaren Rapid liquid capsule, these symptoms may be severe.

Do not take Voltaren Rapid liquid capsule if you:

- currently have a stomach or intestinal ulcer, or other stomach disorder
- have bleeding from the stomach or bowel (symptoms of which may include blood in your stools or black stools)
- have impaired kidney or severe liver disease
- have heart failure.

Do not give this medicine to a child under 14 years of age.

There is not enough information on the use of Voltaren Rapid liquid capsule in children under that age.

Tell your doctor or pharmacist if any of the above apply because Voltaren Rapid liquid capsule is not suitable for you in these circumstances.

Do not take Voltaren Rapid liquid capsule during the last 3 months of pregnancy.

Use of this medicine during the last 3 months of pregnancy may affect your baby and may delay labour and birth. Use of this medicine can increase the risk of miscarriage, particularly when taken close to the time of conception.

Do not take Voltaren Rapid liquid capsule after the expiry date printed on the pack.

Do not take Voltaren Rapid liquid capsule if the foil seal over the capsules has been broken or shows signs of tampering.

In that case, return it to your pharmacist.

If you are not sure whether you should start taking Voltaren Rapid liquid capsule, contact your doctor or pharmacist.

Before you start to take it

Tell your pharmacist or doctor if you are allergic to any other medicines, including aspirin or other NSAID medicines, foods, preservatives or dyes.

They will want to know if you are prone to allergies.

Talk to your doctor before taking Voltaren Rapid liquid capsule if you have:

- asthma or any other chronic lung disease that causes difficulty in breathing.

Talk to your doctor or pharmacist before taking Voltaren Rapid liquid capsule if you have or have had any medical conditions, especially the following:

- established disease of the heart or blood vessels (also called cardiovascular disease, including uncontrolled high blood pressure, congestive heart failure, established ischemic heart disease or peripheral arterial disease) or history of blood vessel disease
- significant risk factors for heart disease, such as high blood pressure, abnormally high levels of fat (cholesterol, triglycerides) in your blood, diabetes or if you smoke
- history of ulcers (stomach or intestinal)
- gastrointestinal problems such as stomach ulcer, bleeding or black stools (vomiting blood or bleeding from the back passage)
- stomach discomfort or heartburn after taking pain relievers or anti-inflammatory medicines

- diseases of the bowel or inflammation of the intestinal tract (Crohn's disease) or colon (ulcerative colitis)
- liver or kidney problems
- a rare liver condition called hepatic porphyria
- bleeding disorder or other blood disorders such as anaemia
- diabetes
- dehydration (eg. by sickness, diarrhoea, before or after recent major surgery)
- swollen feet
- allergic disease, wheezing or shortness of breath or nasal polyps.

Do not exceed the recommended dose.

Treatment with anti-inflammatory medicines, such as diclofenac, may be associated with an increased risk of heart attack (myocardial infarction) or stroke. Any risk is more likely with high doses and prolonged treatment. If you have existing heart disease or risk factors for heart disease, you may be at greater risk. It is generally important to take the lowest dose of Voltaren Rapid liquid capsule that relieves your pain and/or swelling and for the shortest time possible in order to keep your risk for cardiovascular side effects as small as possible.

Tell your pharmacist or doctor if you currently feel unwell or have an infection.

Voltaren Rapid liquid capsule may hide some of the signs of an infection (pain, fever, swelling, redness) and you may think, mistakenly, that you are better or that the infection is not serious.

Tell your pharmacist or doctor if you are pregnant, think you may be pregnant or trying to become pregnant.

Unless absolutely necessary and unless advised by your doctor, do not use Voltaren Rapid liquid capsule during the first 6 months of pregnancy. Use of Voltaren Rapid liquid capsule after 20 weeks of pregnancy may lead to kidney problems in your unborn baby. This could reduce the amount of fluid that surrounds the baby which can be harmful. Do not use at all during the last 3 months of pregnancy.

Tell your pharmacist or doctor if you are breastfeeding.

Unless a doctor has told you to, do not use Voltaren Rapid liquid capsule during breastfeeding.

If you have not told your pharmacist or doctor about any of the above, tell them before you take Voltaren Rapid liquid capsule.

Taking other medicines

Tell your pharmacist or doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines that are important to mention include:

- other anti-inflammatory medicines, eg. aspirin, salicylates, or ibuprofen
- lithium or selective serotonin reuptake inhibitors (SSRIs), medicines used to treat depression
- digoxin, a medicine used to treat heart problems
- ACE inhibitors or beta-blockers, medicines used to treat high blood pressure or heart conditions, glaucoma and migraine
- diuretics, also called fluid or water tablets, used to increase the amount of urine
- medicines used to treat diabetes, except insulin
- prednisone, cortisone, or other corticosteroids, medicines used to provide relief for inflamed areas of the body
- warfarin, other blood thinners or anticoagulants, medicines used to prevent blood clotting
- methotrexate, a medicine used to treat arthritis and some cancers
- cyclosporin or tacrolimus, medicines used by patients who have received organ transplants
- trimethoprim, a medicine used to prevent or treat urinary tract infection
- quinolone antibacterials, medicines used to treat infections
- glucocorticoid, medicines used to treat arthritis
- sulfinpyrazone, a medicine used to treat gout
- voriconazole, a medicine used to treat fungal infections
- phenytoin, a medicine used to treat seizures.

These medicines may be affected by Voltaren Rapid liquid capsule or may affect how well it works. You may need to take different amounts of your medicine, or you may need to take different medications.

Your pharmacist and doctor have more information on medicines to be careful with or to avoid while taking Voltaren Rapid liquid capsule.

How to take it

Follow all directions given to you by your pharmacist or doctor carefully.

These instructions may differ from the information contained in this leaflet.

If you do not understand the instructions, ask your pharmacist or doctor for help.

How much to take

Ask your pharmacist or doctor if you are unsure of the correct dose for you.

They will tell you exactly how much to take. Follow the instructions they give you.

Do not take more than the recommended dose.

Take the lowest dose of Voltaren Rapid liquid capsule that relieves your pain and/or swelling.

Stop taking as soon as your symptoms are relieved. Do not take for longer than a few days at a time.

For adults and children aged 14 years and over:

Take an initial dose of 2 capsules when symptoms appear. If needed, continue with 1 or 2 capsules after 4 to 6 hours. Do not take more than 6 capsules in 24 hours.

How to take it

Swallow the liquid capsule with a glass of water.

When to take it

Take the capsules on an empty stomach (at least half an hour before food or two hours after food).

They will work more quickly if you take them on an empty stomach.

If they upset your stomach, take the capsuls with food.

They will still work if you have to take them with food to prevent stomach upset.

How long to take it

Do not take Voltaren Rapid liquid capsule for more than a few days at a time, unless a doctor has told you to.

It is usually taken for a short time only.

If symptoms persist or worsen, see your pharmacist or doctor to ensure that they are not due to a serious illness.

The use of Voltaren Rapid liquid capsule could mask potentially serious infections.

If you take too much (overdose)

Immediately telephone your doctor or Poisons Information Centre (Australia: 13 11 26, New Zealand: 0800 764 766), or go to Accident and Emergency at your nearest hospital if you think you or anyone else may have taken too much Voltaren Rapid liquid capsule.

Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

While you are taking Voltaren Rapid liquid capsule

Things you must do

If you become pregnant while taking Voltaren Rapid liquid capsule, tell your pharmacist or doctor immediately.

They can discuss with you the risks of using it while you are pregnant.

If, at any time while taking Voltaren Rapid liquid capsule you experience any signs or symptoms of problems with your heart or blood vessels such as chest pain, shortness of breath, weakness or slurring of speech, contact your doctor immediately

If you are about to start taking any new medicine, tell your doctor and pharmacist that you are taking Voltaren Rapid liquid capsule.

If you are going to have surgery, tell the surgeon and the anaesthetist that you are taking Voltaren Rapid liquid capsule.

Anti-inflammatory medicines can slow down blod clotting and affect kidney function.

If you get an infection while taking Voltaren Rapid liquid capsule, tell your doctor.

This medicine may hide some of the signs of an infection (pain, fever, swelling, redness). You may think, mistakenly, that you are better or that the infection is not serious.

Tell all of the doctors, dentists and pharmacists that are treating you that you are taking Voltaren Rapid liquid capsule.

Things you must not do

Do not take any of the following medicines while you are taking Voltaren Rapid liquid capsule without first telling your doctor or pharmacist:

- aspirin or other salicylates
- other medicines containing diclofenac
- any other anti-inflammatory medicine.

If you take these medicines together with Voltaren Rapid liquid capsule, they may cause unwanted side effects.

Do not use Voltaren Rapid liquid capsule to treat any other complaints unless your pharmacist or doctor tells you to.

Do not give your medicine to anyone else, even if they seem to have the same condition as you.

Things to be careful of

Be careful driving, operating machinery or doing jobs that require tbe alert until you know how Voltaren Rapid liquid capsule affects you.

As with other anti-inflammatory medicines, Voltaren Rapid liquid capsule may cause dizziness, drowsiness, spinning sensation (vertigo) or blurred vision in some people. If you have any of these symptoms, do not drive or carry out other activities that need careful attention.

Be careful if you are older than 65 years of age.

Older people may be more sensitive to the effects of Voltaren Rapid liquid capsule than other adults. Take the lowest number of capsules that provides relief of symptoms, and report side effects promptly to your doctor or pharmacist.

Side effects

Tell your pharmacist or doctor as soon as possible if you do not feel well while you are taking Voltaren Rapid liquid capsule.

This medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

If you get an allergic reaction, stop taking and see your doctor immediately.

If you are over 65 years old, you should be especially careful while taking this medicine. Report any side effects promptly to your pharmacist or doctor.

As people get older, they are more likely to get side effects from medicines.

Do not be alarmed by this list of possible side effects. You may not experience any of them. Ask your pharmacist or doctor to answer any questions you may have.

Tell your pharmacist or doctor if you notice any of the following and they worry you:

- stomach upset including nausea (feeling sick), vomiting, heartburn, indigestion, cramps, loss of appetite, wind
- heartburn or pain behind or below the breastbone (possible symptoms of an ulcer in the tube that carries food from the throat to stomach)
- stomach or abdominal pain
- constipation, diarrhoea
- sore mouth or tongue
- altered taste sensation
- headache

- dizziness, spinning sensation
- drowsiness, disorientation, forgetfulness
- feeling depressed, anxious or irritable
- strange or disturbing thoughts or moods
- shakiness, sleeplessness, nightmares
- tingling or numbness of the hands or feet
- feeling of fast or irregular heartbeat, also called palpitations
- unusual weight gain or mild swelling
- symptoms of sunburn (such as redness, itching, swelling, blistering of the lips, eyes, mouth and/or skin) that happen more quickly than normal
- vision disorders (eg. blurred or double vision)
- buzzing or ringing in the ears, difficulty hearing
- hypertension (high blood pressure)
- hair loss or thinning.

Tell your doctor or pharmacist immediately or go to Accident and Emergency at your nearest hospital, if you notice any of the following:

- red or purple skin (possibly signs of blood vessel inflammation)
- severe pain or tenderness in the stomach, vomiting blood or material that looks like coffee grounds, bleeding from the back passage, black sticky bowel motions (stools) or bloody diarrhoea (possible stomach problems)
- sudden or severe itching, rash, skin rash with blisters, itching or hives on the skin, swelling of the face, lips, mouth, tongue, throat or other part of the body which may cause difficulty to swallow, low blood pressure (hypotension), fainting, shortness of breath (possible allergic reaction)
- wheezing, coughing, sudden difficulty in breathing or feelings of tightness in the chest (signs of asthma)]
- yellowing of the skin and/or eyes (signs of hepatitis/liver failure)
- persistent nausea, loss of appetite, unusual tiredness, vomiting, pain in the upper right abdomen, dark urine or pale bowel motions (possible liver problems)
- constant "flu-like" symptoms including chills, fever, sore throat, aching joints, swollen glands, tiredness or lack of energy, bleeding or bruising more easily than normal (possible blood problem)
- painful red areas, large blisters, peeling of layers of skin, bleeding in the lips, eyes, mouth, nose or genitals, which may be accompanied by fever and chills, aching muscles and feeling generally unwell, swollen lymph nodes, facial swelling, blood abnormalities and inflammation of the liver, kidneys and other internal organs (possible serious skin reaction)
- signs of possible effects on the brain, such as sudden and severe headache, stiff neck (signs of viral meningitis), severe nausea, dizziness, numbness, difficulty speaking, paralysis (signs of cerebral attack), convulsion (fits)
- a change in the colour or amount of urine passed, frequent need to urinate, burning feeling when passing urine, blood or excess of protein in the urine (possible kidney disorder)
- sudden and oppressive chest pain (which may be a sign of myocardial infarction or a heart attack)
- breathlessness, difficulty breathing when lying down, swelling of the feet or legs (signs of cardiac failure).

These are very serious side effects. Some of these side effects are signs of a serious allergic reaction. You may need urgent medical attention or hospitalisation. Serious side effects are rare.

Tell your pharmacist or doctor if you notice anything else that is making you feel unwell.

Other side effects not listed above may happen in some people.

After using it

Storage

Keep your capsules in the original packaging until it is time to take them.

If you take the capsules out of the box or blister pack, they may not keep well.

Store the capsules in a cool dry place where the temperature stays below 30°C.

Do not store Voltaren Rapid liquid capsule or any other medicine in the bathroom or near a sink. Do not leave it in the car or on window-sills.

Heat and dampness can destroy some medicines.

Keep the medicine where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your pharmacist or doctor tells you to stop taking Voltaren Rapid liquid capsule or the capsules have passed the expiry date, ask your pharmacist what to do with any medicine that are left over.

Return any unused medicine to your pharmacist for safe disposal.

Product description

What it looks like

Voltaren Rapid liquid capsule are small, yellow liquid capsules. It is available in blister packs of 20.

Ingredients

Voltaren Rapid liquid capsule contains diclofenac potassium as the active ingredient. Each Voltaren Rapid liquid capsule contains 12.5 mg of diclofenac potassium.

The other ingredients are: macrogol 600, gelatin, glycerol, sorbitol, sorbitan, mannitol, and quinoline yellow (E104). Voltaren Rapid liquid capsule also contains trace amounts of medium chain triglycerides and phospholipid based surfactant (Phosal 53 MCT).

Haleon

Sydney, NSW, Australia & Auckland, New Zealand

 **AU: 1800 028 533**
NZ: 0800 540 144

Trade Marks are owned by or licensed to the Haleon group of companies. ©2024 Haleon group of companies or its licensor.

Australian Registration Number

AUST R 171307
AU-NZ 10938 – 10938