

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Advil® Cold & Sinus Liqui-Gels®

**Ibuprofen (free acid and potassium salt)
and Pseudoephedrine Hydrochloride Liqui-Gels**

Read this carefully before you start taking **Advil Cold & Sinus Liqui-Gels** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **Advil Cold & Sinus Liqui-Gels**.

Serious Warnings and Precautions

- If you have glaucoma or difficulty in urination due to an enlarged prostate, do not take this drug unless directed by a doctor.
- Use with caution in those with heart failure, high blood pressure or other conditions that may cause excess fluid collecting in tissues.
- Use with caution in patients at risk of gastrointestinal tract irritation, including those with a history of peptic ulcer. Symptoms may include feeling faint, vomiting blood, bloody or black stools. The chance of stomach bleeding is higher if you are: age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner or steroid drug, take with other drugs containing an NSAID like acetylsalicylic acid (ASA), ibuprofen, naproxen, or prescription anti-inflammatory drugs, or have 3 or more alcoholic drinks every day while using this product.
- Use with caution in patients at risk of kidney problems, such as those with heart failure, liver disease, the elderly or those using diuretics.
- Advil Cold & Sinus Liqui-Gels should not be used during the third trimester of pregnancy.
- Use during the first and second trimesters of pregnancy or nursing should be avoided unless directed by your health care professional.
- Stop use immediately if you observe blood in the urine or you develop urgent, frequent, painful or difficult urination.

What is Advil Cold & Sinus Liqui-Gels used for?

- Fast, effective temporary relief of the symptoms of colds including sore throat pain, sinus pain, nasal congestion, headache, fever, body aches and pains.

How does Advil Cold & Sinus Liqui-Gels work?

Ibuprofen reduces pain and fever. Pseudoephedrine hydrochloride is a nasal decongestant.

What are the ingredients in Advil Cold & Sinus Liqui-Gels?

Medicinal ingredients: ibuprofen (present as free acid and potassium salt) and pseudoephedrine hydrochloride.

Non-medicinal ingredients: D&C yellow no. 10, FD&C red no. 40, fractionated coconut oil, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol.

Advil Cold & Sinus Liqui-Gels comes in the following dosage forms:

Each Liqui-Gel contains ibuprofen 200 mg (present as free acid and potassium salt) and pseudoephedrine hydrochloride 30 mg.

Do not use Advil Cold & Sinus Liqui-Gels if you have or are:

- allergic/hypersensitive to acetylsalicylic acid (ASA), ibuprofen, other salicylates, other NSAIDs, pseudoephedrine or other sympathomimetic amines or any of the ingredients in Advil Cold & Sinus Liqui-Gels (Refer to the nonmedicinal ingredients section of this insert)
- active or recurrent stomach ulcer or gastrointestinal (GI) bleeding or active inflammatory bowel disease (e.g. Crohn's, colitis)
- taking a monoamine oxidase inhibitor (MAOI; e.g. drugs for depression or Parkinson's disease) or for 14 days after stopping the MAOI drug, ASA or other NSAIDs including any other ibuprofen product
- nasal polyps (swelling of the inside of the nose)
- asthma
- allergic manifestations such as anaphylaxis (sudden severe life-threatening allergic reaction), urticaria/hives, rhinitis (stuffed or runny nose that may be due to allergies), skin rash or other allergic symptoms
- dehydrated (significant fluid loss) due to vomiting, diarrhea or lack of fluid intake
- been diagnosed with severe high blood pressure or have heart disease
- right before or after heart surgery
- serious liver disease
- severe kidney disease
- thyroid disease
- Raynaud's Syndrome (a disorder of the circulatory system)
- Systemic Lupus Erythematosus
- in your third trimester of pregnancy

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Advil Cold & Sinus Liqui-Gels. Talk about any health conditions or problems you may have, including if you have or are:

- blood clotting disorder (such as hemophilia)
- breathing problems or chronic lung disease (such as chronic bronchitis)
- diabetes
- difficulty in urination due to prostate enlargement
- glaucoma
- high blood pressure
- mild to moderate kidney disease
- mild to moderate liver disease
- any other serious disease
- under doctor's care for any serious condition
- trying to conceive, in your first or second trimester of pregnancy or if you are breastfeeding
- in the second trimester of pregnancy, your use of NSAIDs, like Advil products, may need to be monitored by a health care practitioner due to the rare risk of kidney problems in the unborn baby which may result in decreased amniotic fluid volume and other complications
- taking medication for weight control
- taking any other drug including over the counter drugs

Other warnings you should know about:

Do not smoke or drink alcohol while using this product.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines. The following may interact with Advil Cold & Sinus Liqui-Gels:

- acetaminophen
- acetylsalicylic acid (ASA)
- allergy medications
- anticoagulants (blood thinning medications)
- selective serotonin reuptake inhibitors (SSRIs) used to treat depression
- anti-hypertensives (blood pressure medications)
- antibiotics (levofloxacin)
- benzodiazepines
- cold medications
- corticosteroids
- cyclosporine
- diabetes medication (including insulin and oral antidiabetic agents)
- digoxin
- diuretics (water pills)
- lithium
- methotrexate
- monoamine

oxidase inhibitors

- nonsteroidal anti-inflammatory drugs (NSAIDs); including naproxen and ibuprofen
- phenytoin
- probenecid
- thyroxine

How to take Advil Cold & Sinus Liqui-Gels:

Usual dose:

- Adults and children 12 to 65 years: Take 1 or 2 Liqui-Gels every 4- 6 hours as needed. Do not exceed 6 Liqui-Gels in 24 hours, unless directed by a doctor.
- Do not give to children under 12, unless directed by a doctor.

Overdose:

If you think you, or a person you are caring for, have taken too much Advil Cold & Sinus Liqui-Gels, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

Continue to take 1 or 2 Liqui-Gels every 4-6 hours as needed after a missed dose. Do not take twice the recommended dose following a missed dose.

What are possible side effects from using Advil Cold & Sinus Liqui-Gels?

Stop use and ask a doctor if:

- you show signs of stomach bleeding
- sore throat pain lasts more than 2 days
- symptoms worsen or last more than 5 days
- fever lasts more than 3 days
- you are nervous, dizzy or can't sleep
- any new symptoms appear

These are not all the possible side effects you may have when taking Advil Cold & Sinus Liqui-Gels. If you experience any side effects not listed here, tell your healthcare professional.

Take with food or milk if upset stomach occurs.

Advil Cold & Sinus Liqui-Gels may occasionally produce unwanted side effects, such as heartburn, constipation, nausea, bloating, nervousness or sleeplessness. Stop use and contact a healthcare professional if these symptoms worsen or persist.

The risk of having side effects may be decreased by using the smallest dose for the shortest duration of time.

Serious side effects and what to do about them

Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Sudden severe headache accompanied by confusion, seizures, vomiting sensation or changes in vision. These may be signs of a serious condition.			✓
Symptoms of allergic reaction, including rash, severe itching/redness, blisters, swelling, or trouble breathing			✓
Blood in vomit, bloody or black stools			✓
Abdominal pain, vomiting, diarrhea		✓	
Ringing or buzzing in the ears / dizziness		✓	
Change in vision		✓	
Fluid retention		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Store at room temperature (15 to 30°C).

Keep out of reach and sight of children. This package contains enough medicine to seriously harm a child.

If you want more information about Advil Cold & Sinus Liqui-Gels:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website: (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); or by contacting the sponsor at www.advil.ca or by calling 1-888-788-8181.

Haleon Canada ULC, Mississauga, ON L5R 4B2

Liqui-Gels® is a trademark or registered trademark of Catalent Pharma Solutions.

This leaflet was prepared by Haleon Canada ULC.

Date of Authorization: 2025-09-26 6200000222491 6200000222492