

LOT  
EXP.



Drug Facts		Informations	
<b>Active ingredients</b> Acetaminophen 325 mg Pseudoephedrine HCl 30 mg	<b>Purposes</b> • Temporary relief of pain • Temporary relief of nasal congestion	<b>Informations générales</b> • Ce médicament est un médicament à usage oral. • Contient du pseudoéphédrine, un médicament à usage oral.	<b>Utilité</b> • Soulage temporairement les symptômes de la grippe et du rhume. • Soulage temporairement les symptômes de la congestion nasale.
<b>Warnings</b> • Do not take if you are allergic to acetaminophen or pseudoephedrine. • Do not take if you have ever had a seizure. • Do not take if you have ever had a heart attack or stroke. • Do not take if you have ever had a blood clot. • Do not take if you have ever had a kidney problem. • Do not take if you have ever had a liver problem. • Do not take if you have ever had a stomach or intestinal problem. • Do not take if you have ever had a urinary tract infection. • Do not take if you have ever had a thyroid problem. • Do not take if you have ever had a diabetes problem. • Do not take if you have ever had a high blood pressure problem. • Do not take if you have ever had a low blood pressure problem. • Do not take if you have ever had a heart rhythm problem. • Do not take if you have ever had a blood sugar problem. • Do not take if you have ever had a blood cholesterol problem. • Do not take if you have ever had a blood calcium problem. • Do not take if you have ever had a blood iron problem. • Do not take if you have ever had a blood magnesium problem. • Do not take if you have ever had a blood potassium problem. • Do not take if you have ever had a blood sodium problem. • Do not take if you have ever had a blood zinc problem.	<b>Contra-indications</b> • Ne prenez pas si vous êtes allergique à l'acétaminophène ou au pseudoéphédrine. • Ne prenez pas si vous avez déjà eu une crise d'épilepsie. • Ne prenez pas si vous avez déjà eu un infarctus du myocarde ou un AVC. • Ne prenez pas si vous avez déjà eu un caillot sanguin. • Ne prenez pas si vous avez déjà eu un problème rénal. • Ne prenez pas si vous avez déjà eu un problème hépatique. • Ne prenez pas si vous avez déjà eu un problème stomacal ou intestinal. • Ne prenez pas si vous avez déjà eu une infection des voies urinaires. • Ne prenez pas si vous avez déjà eu un problème thyroïdien. • Ne prenez pas si vous avez déjà eu un problème diabétique. • Ne prenez pas si vous avez déjà eu un problème de haute ou basse tension artérielle. • Ne prenez pas si vous avez déjà eu un problème de rythme cardiaque. • Ne prenez pas si vous avez déjà eu un problème de glycémie. • Ne prenez pas si vous avez déjà eu un problème de cholestérol. • Ne prenez pas si vous avez déjà eu un problème de calcium sanguin. • Ne prenez pas si vous avez déjà eu un problème de fer sanguin. • Ne prenez pas si vous avez déjà eu un problème de magnésium sanguin. • Ne prenez pas si vous avez déjà eu un problème de potassium sanguin. • Ne prenez pas si vous avez déjà eu un problème de sodium sanguin. • Ne prenez pas si vous avez déjà eu un problème de zinc sanguin.		
<b>Directions</b> • Adults and children 12 years and over: Take 2 tablets every 4 to 6 hours. Do not take more than 6 tablets in 24 hours. Do not take if you have ever had a seizure.	<b>Mode d'emploi</b> • Prenez 2 comprimés toutes les 4 à 6 heures. Ne prenez pas plus de 6 comprimés en 24 heures. Ne prenez pas si vous avez déjà eu une crise d'épilepsie.	<b>Autres renseignements</b> • Ce médicament est un médicament à usage oral. • Contient du pseudoéphédrine, un médicament à usage oral.	<b>Autres renseignements</b> • Ce médicament est un médicament à usage oral. • Contient du pseudoéphédrine, un médicament à usage oral.

**NEOCITRAN**  
EXTRA STRENGTH COLD & CONGESTION FORT  
Lemon / Citron

10 SINGLE DOSE POUCHES (POUCHES)  
10 SACHETS À DOSE UNIBLÉ (SACHETS)

127.5461531

0 58478 03703 3

Contient l'acétaminophène et d'autres ingrédients / Contient de l'acétaminophène et d'autres ingrédients

**HALEON**

**NEOCITRAN**  
EXTRA STRENGTH COLD & CONGESTION FORT  
Lemon / Citron

**EXTRA STRENGTH COLD & CONGESTION FORT**  
RHYME ET CONGESTION

**Sinus & nasal congestion / Congestion nasale et sinusale**  
**Sore throat and headache / Maux de gorge et maux de tête**

10 SINGLE DOSE POUCHES (POUCHES)  
10 SACHETS À DOSE UNIBLÉ (SACHETS)

**Drug Facts (continued)**

**Directions**  
• Adults and children 12 years and over: Take 2 tablets every 4 to 6 hours. Do not take more than 6 tablets in 24 hours. Do not take if you have ever had a seizure.

**Other information**  
• Do not take if you are allergic to acetaminophen or pseudoephedrine.  
• Do not take if you have ever had a seizure.  
• Do not take if you have ever had a heart attack or stroke.  
• Do not take if you have ever had a blood clot.  
• Do not take if you have ever had a kidney problem.  
• Do not take if you have ever had a liver problem.  
• Do not take if you have ever had a stomach or intestinal problem.  
• Do not take if you have ever had a urinary tract infection.  
• Do not take if you have ever had a thyroid problem.  
• Do not take if you have ever had a diabetes problem.  
• Do not take if you have ever had a high blood pressure problem.  
• Do not take if you have ever had a low blood pressure problem.  
• Do not take if you have ever had a heart rhythm problem.  
• Do not take if you have ever had a blood sugar problem.  
• Do not take if you have ever had a blood cholesterol problem.  
• Do not take if you have ever had a blood calcium problem.  
• Do not take if you have ever had a blood iron problem.  
• Do not take if you have ever had a blood magnesium problem.  
• Do not take if you have ever had a blood potassium problem.  
• Do not take if you have ever had a blood sodium problem.  
• Do not take if you have ever had a blood zinc problem.

**Active ingredients**  
Acetaminophen 325 mg  
Pseudoephedrine HCl 30 mg

**Other information**  
• Do not take if you are allergic to acetaminophen or pseudoephedrine.  
• Do not take if you have ever had a seizure.  
• Do not take if you have ever had a heart attack or stroke.  
• Do not take if you have ever had a blood clot.  
• Do not take if you have ever had a kidney problem.  
• Do not take if you have ever had a liver problem.  
• Do not take if you have ever had a stomach or intestinal problem.  
• Do not take if you have ever had a urinary tract infection.  
• Do not take if you have ever had a thyroid problem.  
• Do not take if you have ever had a diabetes problem.  
• Do not take if you have ever had a high blood pressure problem.  
• Do not take if you have ever had a low blood pressure problem.  
• Do not take if you have ever had a heart rhythm problem.  
• Do not take if you have ever had a blood sugar problem.  
• Do not take if you have ever had a blood cholesterol problem.  
• Do not take if you have ever had a blood calcium problem.  
• Do not take if you have ever had a blood iron problem.  
• Do not take if you have ever had a blood magnesium problem.  
• Do not take if you have ever had a blood potassium problem.  
• Do not take if you have ever had a blood sodium problem.  
• Do not take if you have ever had a blood zinc problem.

**Warnings**  
• Do not take if you are allergic to acetaminophen or pseudoephedrine.  
• Do not take if you have ever had a seizure.  
• Do not take if you have ever had a heart attack or stroke.  
• Do not take if you have ever had a blood clot.  
• Do not take if you have ever had a kidney problem.  
• Do not take if you have ever had a liver problem.  
• Do not take if you have ever had a stomach or intestinal problem.  
• Do not take if you have ever had a urinary tract infection.  
• Do not take if you have ever had a thyroid problem.  
• Do not take if you have ever had a diabetes problem.  
• Do not take if you have ever had a high blood pressure problem.  
• Do not take if you have ever had a low blood pressure problem.  
• Do not take if you have ever had a heart rhythm problem.  
• Do not take if you have ever had a blood sugar problem.  
• Do not take if you have ever had a blood cholesterol problem.  
• Do not take if you have ever had a blood calcium problem.  
• Do not take if you have ever had a blood iron problem.  
• Do not take if you have ever had a blood magnesium problem.  
• Do not take if you have ever had a blood potassium problem.  
• Do not take if you have ever had a blood sodium problem.  
• Do not take if you have ever had a blood zinc problem.

DE30604CA-13

LOT  
EXP.

Drug Facts	Indicaciones	Purposes	Información	Usos
<b>Active ingredients</b> Pseudoephedrine HCl 120 mg Acetaminophen 325 mg Dextromethorphan HBr 10 mg	<b>Relieves</b> nasal congestion, sinus pain, sore throat, cough, and cold/flu symptoms.	<b>Relieves</b> pain and fever. <b>Reduces</b> cough. <b>Relieves</b> cold/flu symptoms.	<b>Indicaciones:</b> alivia la congestión nasal, el dolor de senos, el dolor de garganta, la tos y los síntomas de resaca y gripe.	<b>Usos:</b> alivia temporalmente los síntomas de congestión nasal, dolor de senos, dolor de garganta, tos y síntomas de resaca y gripe.
<b>Warnings:</b> Acetaminophen may cause severe or possibly fatal liver damage if you take too much or if you take it with other drugs containing acetaminophen. Do not drink alcohol while taking this medicine. Do not take more than the directed amount. Do not take this medicine if you are taking other medicines that contain acetaminophen. Do not take this medicine if you are taking other medicines that contain pseudoephedrine. Do not take this medicine if you are taking other medicines that contain dextromethorphan. Do not take this medicine if you are taking other medicines that contain aspirin. Do not take this medicine if you are taking other medicines that contain ibuprofen. Do not take this medicine if you are taking other medicines that contain naproxen. Do not take this medicine if you are taking other medicines that contain tramadol. Do not take this medicine if you are taking other medicines that contain tramadol.	<b>Precauciones:</b> Evite el alcohol. No tome más de la dosis recomendada. No tome este medicamento si está tomando otros medicamentos que contengan acetaminofeno, pseudoefedrina o dextrometorfano. No tome este medicamento si está tomando otros medicamentos que contengan aspirina, ibuprofeno o naproxeno. No tome este medicamento si está tomando otros medicamentos que contengan tramadol.	<b>Warnings:</b> Acetaminophen may cause severe or possibly fatal liver damage if you take too much or if you take it with other drugs containing acetaminophen. Do not drink alcohol while taking this medicine. Do not take more than the directed amount. Do not take this medicine if you are taking other medicines that contain acetaminophen. Do not take this medicine if you are taking other medicines that contain pseudoephedrine. Do not take this medicine if you are taking other medicines that contain dextromethorphan. Do not take this medicine if you are taking other medicines that contain aspirin. Do not take this medicine if you are taking other medicines that contain ibuprofen. Do not take this medicine if you are taking other medicines that contain naproxen. Do not take this medicine if you are taking other medicines that contain tramadol. Do not take this medicine if you are taking other medicines that contain tramadol.	<b>Contraindicaciones:</b> No tome este medicamento si está tomando otros medicamentos que contengan acetaminofeno, pseudoefedrina o dextrometorfano. No tome este medicamento si está tomando otros medicamentos que contengan aspirina, ibuprofeno o naproxeno. No tome este medicamento si está tomando otros medicamentos que contengan tramadol.	<b>Contraindicaciones:</b> No tome este medicamento si está tomando otros medicamentos que contengan acetaminofeno, pseudoefedrina o dextrometorfano. No tome este medicamento si está tomando otros medicamentos que contengan aspirina, ibuprofeno o naproxeno. No tome este medicamento si está tomando otros medicamentos que contengan tramadol.

Indicaciones	Modo de empleo	Ingredientes
<b>Indicaciones:</b> alivia la congestión nasal, el dolor de senos, el dolor de garganta, la tos y los síntomas de resaca y gripe.	<b>Modo de empleo:</b> Tómelo con un vaso de agua. Tómelo cada 4 horas, pero no más de 6 veces al día. No tome más de 1200 mg de pseudoefedrina o 4000 mg de acetaminofeno al día. No tome más de 10 mg de dextrometorfano al día. No tome este medicamento si está tomando otros medicamentos que contengan acetaminofeno, pseudoefedrina o dextrometorfano. No tome este medicamento si está tomando otros medicamentos que contengan aspirina, ibuprofeno o naproxeno. No tome este medicamento si está tomando otros medicamentos que contengan tramadol.	<b>Ingredientes:</b> Pseudoefedrina HCl, Acetaminofeno, Dextrometorfano HBr, Citrato de sodio, Difenidramina HCl, Mentol, Aceite de mentol, Aceite de eucalipto, Aceite de mentol, Aceite de eucalipto, Aceite de mentol, Aceite de eucalipto.

**NEOCITRAN**

EXTRA STRENGTH EXTRA RHYME ET CONGESTION FORT

DAY / NIGHT

Contra le rhume et la congestion / Contiene el acetaminofeno y otros ingredientes

HALLEON

Lemon / Citron

10 SINGLE DOSE POUCHES (POUCHES) / 10 SAQUETS A DOSER (INDIVIDUALES)

Drug Facts (continued)
<b>Directions:</b> Adults: Take 1 pouch every 4 hours, with or without food. Do not take more than 10 pouches in 24 hours. Children under 12: Do not take. Do not take if you are pregnant or breastfeeding. Do not take if you are taking other medicines that contain acetaminophen, pseudoephedrine, or dextromethorphan. Do not take if you are taking other medicines that contain aspirin, ibuprofen, or naproxen. Do not take if you are taking other medicines that contain tramadol. Do not take if you are taking other medicines that contain tramadol.
<b>Other information:</b> Keep out of reach of children. Do not use if the pouches are torn, damaged, or missing. Do not use if the pouches are past their expiration date. Do not use if the pouches are past their expiration date. Do not use if the pouches are past their expiration date.
<b>Active ingredients:</b> Pseudoephedrine HCl, Acetaminophen, Dextromethorphan HBr, Citrate de sodium, Difenidramine HCl, Mentol, Aceite de mentol, Aceite de eucalipto, Aceite de mentol, Aceite de eucalipto.
<b>Warnings:</b> Acetaminophen may cause severe or possibly fatal liver damage if you take too much or if you take it with other drugs containing acetaminophen. Do not drink alcohol while taking this medicine. Do not take more than the directed amount. Do not take this medicine if you are taking other medicines that contain acetaminophen. Do not take this medicine if you are taking other medicines that contain pseudoephedrine. Do not take this medicine if you are taking other medicines that contain dextromethorphan. Do not take this medicine if you are taking other medicines that contain aspirin. Do not take this medicine if you are taking other medicines that contain ibuprofen. Do not take this medicine if you are taking other medicines that contain naproxen. Do not take this medicine if you are taking other medicines that contain tramadol. Do not take this medicine if you are taking other medicines that contain tramadol.
<b>Directions:</b> Adults: Take 1 pouch every 4 hours, with or without food. Do not take more than 10 pouches in 24 hours. Children under 12: Do not take. Do not take if you are pregnant or breastfeeding. Do not take if you are taking other medicines that contain acetaminophen, pseudoephedrine, or dextromethorphan. Do not take if you are taking other medicines that contain aspirin, ibuprofen, or naproxen. Do not take if you are taking other medicines that contain tramadol. Do not take if you are taking other medicines that contain tramadol.
<b>Other information:</b> Keep out of reach of children. Do not use if the pouches are torn, damaged, or missing. Do not use if the pouches are past their expiration date. Do not use if the pouches are past their expiration date. Do not use if the pouches are past their expiration date.
<b>Active ingredients:</b> Pseudoephedrine HCl, Acetaminophen, Dextromethorphan HBr, Citrate de sodium, Difenidramine HCl, Mentol, Aceite de mentol, Aceite de eucalipto, Aceite de mentol, Aceite de eucalipto.
<b>Warnings:</b> Acetaminophen may cause severe or possibly fatal liver damage if you take too much or if you take it with other drugs containing acetaminophen. Do not drink alcohol while taking this medicine. Do not take more than the directed amount. Do not take this medicine if you are taking other medicines that contain acetaminophen. Do not take this medicine if you are taking other medicines that contain pseudoephedrine. Do not take this medicine if you are taking other medicines that contain dextromethorphan. Do not take this medicine if you are taking other medicines that contain aspirin. Do not take this medicine if you are taking other medicines that contain ibuprofen. Do not take this medicine if you are taking other medicines that contain naproxen. Do not take this medicine if you are taking other medicines that contain tramadol. Do not take this medicine if you are taking other medicines that contain tramadol.

0 58478 03703 3

127.5M45131

DE30904CA-13