

# PANADOL EXTRA-STRENGTH

## Product Information

Each tablet of **Panadol Extra-Strength** contains:

**Active ingredient** Acetaminophen 500mg

**Purpose:** analgesic/antipyretic. Excipients q.e. 1 tablet.

### Use Panadol Extra Strength For:

Relief of mild to moderate pain including:

Headache.

Muscle pain.

Osteoarthritis pain.

Fever and pain after vaccination.

Pain after dental procedures / tooth extraction.

In addition, it reduces fever.

## Dosage

**Adults (12 years and older):** Oral administration 2 tablets every 6 hours. Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period necessary. Do not take more frequently than every 4 hours. Maximum 8 tablets per 24 hours. Not recommended in children under 12 years. Do not administer more than 4 times in a day the recommended dose, or for more than 10 consecutive days to adults. Do not exceed the stated dose.

## Contraindications

Do not use if you are allergic to acetaminophen, or any other ingredients in the product.

## Warnings and precautions

Contains acetaminophen. Do not use this medicine if you are taking any other medicines containing acetaminophen to treat pain, fever, symptoms of cold and flu or to aid sleep. Do not take more than recommended dose as it may cause serious harm to your liver. Always read and follow the label. **Check with your doctor before use if you:** Have liver or kidney problems, are taking any medication or if you are under medical treatment, you are pregnant, have a severe infection, are severely underweight, or are chronic heavy alcohol user, you may need to avoid using this product altogether or limit the amount of acetaminophen that you take, as this may increase the risk of metabolic acidosis, signs of metabolic acidosis include: deep, rapid, difficult breathing, nausea, vomiting, loss of appetite. Contact a doctor immediately if you get a combination of these symptoms. Pain or fever persists for more than 3 days, gets worse or new symptoms occur. You can take this product whilst breastfeeding. Consider taking the lowest effective dose for the shortest period of time. Keep out of sight and reach of children.

## Drug interactions

Consult your doctor if you are taking warfarin or similar medicines.

## Adverse reactions

**Stop use if:** You have skin allergic reactions, breathing problems, swelling of lips, tongue, face or throat, peeling or mouth ulcers; unexplained bruising or bleeding. You experience skin reactions such as Stevens-Johnson Syndrome, Toxic Epidermal Necrolysis or Acute Generalized Exanthematous Pustulosis. You have previously experienced breathing problems with acetylsalicylic acid or non-steroidal anti-inflammatories and experience a similar reaction. These reactions are rare.

## Overdose

**If you exceed the recommended dose:** Seek medical advice from a doctor immediately even if you do not have any symptoms because of the risk of liver failure.

## Other information

Store below 30°C. Oral use. Free Sale. Use only if pouch is intact. To report adverse events [SafetyReportingAmericas@haleon.com](mailto:SafetyReportingAmericas@haleon.com), for quality complaints [PQC.caricam@haleon.com](mailto:PQC.caricam@haleon.com) or by phone: Jamaica: +18766302074, Trinidad & Tobago: +18682245759. [www.haleon.com](http://www.haleon.com). [www.panadol.com](http://www.panadol.com)