



Children's Hayfever & Allergy 1mg/ml syrup

Cetirizine Hydrochloride

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your pharmacist has told you.

This medicine is available without prescription, but you still need to use Piriteze Children's Hayfever & Allergy Syrup carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse.

In this leaflet:

1. What Piriteze Children's Hayfever & Allergy Syrup is and what it is used for
2. What you need to know before you take Piriteze Children's Hayfever & Allergy Syrup
3. How to take Piriteze Children's Hayfever & Allergy Syrup
4. Possible side effects
5. How to store Piriteze Children's Hayfever & Allergy Syrup
6. Contents of the pack and other information

1. What Piriteze Children's Hayfever & Allergy Syrup is and what it is used for

Piriteze Children's Hayfever & Allergy Syrup is used to treat the allergic symptoms of hay fever (itchy or watery eyes, sneezing and a runny nose) and other allergies e.g. pet, house dust mite and mould spore allergies. It can also be used to relieve the rashes and itching of urticaria (hives). The active ingredient is cetirizine hydrochloride, an antihistamine which can help to relieve the symptoms of some allergies and itchy skin rashes.

2. What you need to know before you take Piriteze Children's Hayfever & Allergy Syrup



Do not take Piriteze Children's Hayfever & Allergy Syrup

- If you have ever had an allergic reaction to antihistamines or to any of the other ingredients (listed in Section 6)
- If you have severe kidney problems



Take special care with Piriteze Children's Hayfever & Allergy Syrup

- If you have an intolerance to some sugars, contact your doctor before taking this medicine.
- Be careful when drinking alcohol while using Piriteze Children's Hayfever & Allergy Syrup. It

can increase the effects of drinking.

- Do not drive or operate machinery if the syrup makes you feel drowsy. At the recommended dose, the medicine does not cause drowsiness in the majority of people.

- If you have kidney problems as you may need to take a lower dose (see How to take Piriteze Children's Hayfever & Allergy Syrup).

- If you are epileptic or at risk of convulsions (fits).

- If you are having an allergy skin test you need to wait 3 days after taking Piriteze Children's Hayfever & Allergy Syrup.

- If you have an enlarged prostate. It may increase the risk of having difficulty passing water.



If you are taking other medicines

Talk to your doctor or pharmacist before taking this syrup if you are taking any prescribed medicines; including medicines obtained without a prescription.



Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking Piriteze Children's Hayfever & Allergy Syrup if you are pregnant or breast-feeding.





3. How to take Piriteze Children's Hayfever & Allergy Syrup

Adults and adolescents aged 12 years and over: **Take two 5 ml spoonfuls once daily.**

Children aged between 6 and 12 years: **Give one 5 ml spoonful twice daily - morning and evening.**

Children aged between 2 and 6 years: **Give half a spoonful (2.5 ml) twice daily - morning and evening.**

Children under 2 years: **Not recommended.**

For people with moderate kidney problems: **Take one 5 ml spoonful once daily.**



- **Do not take more than the recommended dose.**
- **Do not give to children under 2 years.**

If you or a child take too much syrup

Contact your doctor or casualty department. Do not drive if you have taken too much.

If you forget to take the syrup

Take a dose as soon as you remember, unless it is nearly time to take the next one. Never take two doses together.

If you stop taking the syrup:

- Rarely, pruritus (intense itching) and/or urticaria may occur. If your symptoms persist, **see your doctor.**
- Seek medical advice if withdrawal symptoms are experienced.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Piriteze Children's Hayfever & Allergy Syrup can have side effects, but not everybody gets them.

Stop taking the medicine and seek immediate medical help if you experience:

- Sudden signs of an allergic reaction such as developing a rash, swelling of the mouth or face or difficulty breathing
- Bruising or bleeding more easily than normal
- Convulsions or fits
- Loss of consciousness.

Stop taking the medicine and tell your doctor if you experience:

Confusion, hallucinations, abnormal liver function tests or a rapid heartbeat.

The following side effects may occur:

Common (may affect up to 1 in 10 people): Headache or dizziness, drowsiness or tiredness, restlessness, dry mouth, sore throat, abdominal pain, diarrhoea, nausea [feeling sick], runny nose.

Uncommon (may affect up to 1 in 100 people): Agitation, weakness, malaise, pins and needles, itching, rash or hives.

Rare (may affect up to 1 in 1,000 people): Aggression, depression, weight gain, swelling, insomnia, movement disorders including tremor, muscle spasm and tics.

Very Rare (may affect up to 1 in 10,000 people): Taste disorder, low blood platelets count, blurred vision, eye movement disorder, difficult or painful or involuntary urination.

At a very low frequency that cannot be estimated: Increased appetite, nightmares, inflammation of the liver, hepatitis, skin eruptions, joint pain, memory loss, difficulty passing water or a feeling of dizziness. In very rare cases people have had suicidal thoughts when taking this medicine. If you experience these, stop taking the medicine and see your doctor.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Piriteze Children's Hayfever & Allergy Syrup

Keep out of the sight and reach of children.

Do not use this medicine after the 'EXP' date shown on the pack.

6. Contents of the pack and other information

Active ingredient Each 5 ml dose contains Cetirizine Hydrochloride 5 mg.

Other ingredients Glycerol, propylene glycol, sorbitol (E 420), methyl parahydroxybenzoate (E 218), propyl parahydroxybenzoate (E 216), sodium acetate, acetic acid, saccharin sodium, banana flavour and purified water. **Methyl and propyl parahydroxybenzoates (E 218 and E 216) may cause allergic reactions (possibly delayed).**

The bottle contains 70 ml.

The marketing authorisation holder is Haleon UK Trading Limited, The Heights, Weybridge, KT13 0NY, U.K and all enquiries should be sent to this address.

The manufacturer is Omega Pharma Manufacturing, GmbH & Co. KG, Benzstr. 25, 71083 Herrenberg, Germany.

This leaflet was last revised in February 2026.

Trade Marks are owned by or licensed to the Haleon group of companies.

