Cut along this line to open.

HALEON

# Panaflex

## Cool Patch

***Fast acting, long lasting**
- **Muscle & joint pain**
- **Neck & shoulder ache**
- **Backache**

**5 patches**

**MEDIUM size 8 x 6cm**

---

# Panaflex Cool Patch

*Contains Medicinal Ingredients*
✓ *Sticks firmly on the site of pain yet removes easily* ✓ *Non-greasy* ✓ *Does not stain clothes*
✓ *Contains Vitamin E*

**Indications:** For the temporary relief of muscular and joint pain e.g. due to simple backache, strains, sprains. **Dosage and Administration:** For topical use only. **Maximum duration of use without medical advice: 7 days.** One patch should be applied to the affected area 1 to 2 times a day, as required. **Maximum total daily dose: 2 patches, even if there is more than one injured area to be treated. For use in adults and children aged 12 years and above. Children under 12 years of age: seek medical advice before use.** **Contraindications:** Contraindicated in children below 2 years of age and in patients with known hypersensitivity to glycol salicylate, menthol or tocopherol acetate or any other ingredient of the preparation. **Warnings and Precautions:** Use the product with caution in patients with history of allergic reaction to aspirin or other NSAIDs. The product should be applied only to intact, non-diseased skin and not to skin wounds. The product should not come into contact with the eyes, nostrils, and other mucous membranes. In case of accidental contact, the eyes or other mucous membranes should be rinsed well with water. The product should not be used with external heat, such as electric heating pad, as it may result in excessive skin irritations or skin burns. Some people may experience discomfort, particularly those with sensitive skin, if the product is used after a bath. If the condition does not improve or worsens within 7 days of starting treatment, patients should seek medical advice to exclude an alternative underlying cause of pain. Discontinue the treatment if skin irritation occurs. Not for use in children under 12 years of age unless directed by a healthcare professional. Use with caution in older children. Seizures may occur in children with a history of seizures due to the presence of terpene derivatives (menthol). Keep out of the sight and reach of children. **Interactions:** If you are taking anticoagulants (eg warfarin) and antiplatelet medications, please consult your doctor before using this product as it may result in increased risk of bleeding. **Pregnancy and Lactation:** Do not use during pregnancy or when breastfeeding. **Side Effects:** Application site hypersensitivity, application site reaction, skin burning sensation, rash and pruritus. **Symptoms and Treatment of Overdose:** Overdose after incorrect topical application or due to accidental ingestion for products containing salicylates and menthol can be associated with a number of symptoms which can include: nausea, vomiting, severe abdominal pain, sweating, tinnitus, vertigo, deafness, dizziness, ataxia, headache, confusion, drowsiness and coma. Relatively large amounts of vitamin E usually cause no harm but occasionally muscle weakness, fatigue, nausea, and diarrhoea occur. In the event of overdose, supportive care may be required. Further management should be as clinically indicated or as recommended by the national poisons center, where available. **Storage Condition:** Store below 30°C. **Indikasi:** Memberi kelegaan sementara sakit otot dan sendi pada badan seperti sakit belakang yang sederhana, ketegangan, atau kejadian terseliuh. **Arahan Penggunaan:** Untuk penggunaan topikal sahaja. **Tempoh maksimum penggunaan tanpa nasihat perubatan: 7 hari.** Satu ubat tampalan harus digunakan pada bahagian badan yang sakit 1 atau 2 kali sehari mengikut keperluan. **Dos harian maksimum: 2 ubat tampalan, walaupun terdapat lebih daripada satu bahagian badan yang cedera untuk dirawat. Untuk penggunaan orang dewasa dan kanak-kanak berumur 12 tahun ke atas. Kanak-kanak di bawah 12 tahun: dapatkan nasihat perubatan sebelum guna.** **Kontraindikasi:** Dikontraindikasikan untuk kanak-kanak di bawah umur 2 tahun dan untuk pesakit yang diketahui dengan hipersensitiviti terhadap *glycol salicylate*, mentol atau *tocopherol acetate* atau ramuan lain yang terkandung dalam produk ini. **Amaran dan Perhatian:** Gunakan produk ini dengan berhati-hati untuk pesakit yang mempunyai sejarah reaksi alahan terhadap aspirin atau ubat-ubatan anti radang tanpa steroid (NSAIDs) yang lain. Produk ini hanya boleh digunakan pada kulit yang utuh dan tidak berpenyakit dan bukan pada kulit yang luka. Produk ini tidak harus bersentuhan dengan mata, lubang hidung, dan membran mukus yang lain. Sekiranya terkena secara tidak sengaja, mata atau membran mukus yang lain harus dibilas dengan air. Produk ini tidak boleh digunakan bersama haba luaran, seperti pad pemanasan elektrik, yang mungkin menyebabkan kerengsaan kulit berlebihan atau kulit terbakar. Sesetengah orang mungkin mengalami ketidakselesaan, terutamanya mereka yang mempunyai kulit sensitif, jika produk ini digunakan selepas mandi. Jika keadaan tidak beransur pulih atau bertambah serius dalam masa 7 hari sejak memulakan rawatan, pesakit harus mendapatkan nasihat perubatan untuk mengecualikan penyebab alternatif kesakitan. Hentikan rawatan jika berlaku kerengsaan kulit. Tidak boleh digunakan untuk kanak-kanak di bawah umur 12 tahun kecuali jika diarahkan oleh pakar kesihatan. Gunakan dengan berhati-hati pada kanak-kanak yang lebih tua. Sawan boleh berlaku pada kanak-kanak yang pernah mengalami sawan disebabkan oleh kandungan derivatif *terpene* (mentol). Jauhkan daripada pandangan dan capaian kanak-kanak. **Interaksi:** Jika anda mengambil ubat-ubatan *anticoagulant* (seperti *warfarin*) dan *antiplatelet,* sila dapatkan nasihat doktor anda sebelum menggunakan produk ini sebab kemungkinan peningkatan risiko pendarahan. **Kehamilan dan Penyusuan:** Jangan guna semasa kehamilan atau penyusuan. **Kesan Sampingan:** Hipersensitiviti, reaksi, sensasi terbakar pada kulit, ruam dan *pruritus* di sekitar permukaan ubat tampalan digunakan. **Gejala-gejala dan Rawatan Terlebih Dos:** Terlebih dos ubat selepas penggunaan topikal yang tidak betul atau akibat secara tidak sengaja termakan produk-produk yang mengandungi *salicylates* dan mentol yang diketahui berkaitan dengan beberapa gejala, termasuk: loya, muntah, sakit perut yang teruk, berpeluh, *tinnitus, vertigo*, pekak, pening, *ataxia*, sakit kepala, kekeliruan, mengantuk dan koma. Jumlah vitamin E yang agak besar biasanya tidak mendatangkan kesan mudarat tetapi kadangkala kelemahan otot, keletihan, loya dan cirit-birit berlaku. Jika penggunaan dos yang berlebihan berlaku, rawatan sokongan mungkin diperlukan. Pengurusan lanjut harus seperti yang diindikasikan secara klinikal atau seperti yang disyorkan oleh pusat racun negara, jika ada. **Kaedah Penyimpanan:** Simpan di bawah suhu 30°C. **Active Ingredients:** L-Menthol 4.025% w/w, Glycol Salicylate 1.050% w/w, DL-α-Tocopherol Acetate (Vitamin E) 0.875% w/w. **Inactive Ingredients:** Zinc Oxide, Silicon Dioxide, Aluminium Chlorohydrate, Titanium Dioxide, Adhesive Plaster, Purified Lanolin, Paraffin Oil. Source of Purified Lanolin: ovine. *Refers to overall sensation felt or perceived after 80 seconds for fast acting and overall sensation up to 7.5 hours for long lasting.

***Manufacturer:*** Teh Seng Pharmaceutical Mfg. Co., Ltd. Secondary Factory, No. 42-1, Huan Gong Road, Yong Kang District, Tainan City, Taiwan, R.O.C.

***Product Registration Holder:*** Malaysia - Haleon Malaysia Sdn. Bhd. (Registration No: 195901000115 (3467-X)) Lot 89, Jalan Enggang, Ampang/ Hulu Kelang Industrial Estate, 68000 Ampang, Selangor, Malaysia. ***Imported by:*** Brunei - Kingston Beverage & Creamery Sdn. Bhd. Plot 73, Lot 3, 4, 5 & 6, Beribi Industrial Complex, Jalan Gadong BE1118, Negara Brunei Darussalam. Tel: 673-2450102.
Singapore - Haleon Singapore Pte. Ltd. 23 Rochester Park #03-02, Singapore 139234.

Trade Marks are owned by or licensed to the Haleon group of companies.
©2025 Haleon group of companies or its licensor.

MY: mystory.my@haleon.com
SG: mystory.sg@haleon.com
MY: 1800 88 3225
SG: 1800 622 7238

BATCH NO:
MFG:
EXP:

9 556019 304499

PR-130642MYAD MAL21066034XR BRU24043429NP