100

Advil
Relieves Pain FAST
P4113801
Medicine Information
Active Ingredient (in each liquid capsule)
Ibuprofen 200 mg
Uses
Advil provides temporary relief of pain and discomfort from:
• Headache • Period Pain • Sore Throat Pain
• Back Pain • Dental Pain • Cold & Flu
• Muscle Pain • Arthritis Pain
Advil reduces fever.
Warnings
Do not use this product:
• If you are allergic to ibuprofen, aspirin or other anti-inflammatory medicine, pain reliever / fever reducer or any other component of this product
• If you are pregnant, or trying to become pregnant
• If you have a current or past history of stomach ulcer / bleeding / disorders
• If you have impaired kidney function or heart failure
• In children under 12 years
Unless your doctor has told you do not use this product:
• If you have asthma, high blood pressure, heart or liver disease
• If you are breast feeding
• With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
• If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin
• If you are 65 years or over or under a doctor's care for any serious condition
While using this product:
• Stop use and ask your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
• Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
• Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage
• Side effects may be minimised by using the smallest dose for the shortest duration of time
• In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)
Contains sulfites, soya bean products, and potassium (17,7 mg per capsule)
Directions For Use
For use by Adults and Children from 12 years and over.
Begin by taking 1 or 2 capsules, swallowed whole with water. One capsule will often be adequate for mild pain, but for more severe pain you may require two capsules.
Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 capsules in 24 hours.
Continued...
Advil
Relieves Pain FAST
HALEON
20 Liquid Capsules
Advil
Ibuprofen 200 mg
Relieves Pain FAST
AUST R 97229
20 Liquid Capsules
9 310488 017157
Medicine Information (continued)
Other Information:
STORE BELOW 25°C
IN A DRY PLACE
Do not use this product if the foil is broken
Haleon
Sydney NSW, Australia &
Auckland, New Zealand
AU: 1800 028 533
NZ: 0800 540 144
www.advil.net.au
Trade Marks owned or licensed by Haleon.
©2023 Haleon or licensor.
KEEP OUT OF REACH OF CHILDREN

100

Mfg. Lic. No.: KTK/251/422/99

Advil LIQUID CAPS

Medicine Information

Active Ingredient (in each soft capsule)
Ibuprofen 200 mg

Uses
Temporary relief of pain and discomfort from:
- Headache
- Back Pain
- Muscle Pain
- Period Pain
- Dental Pain
- Arthritis Pain
- Sore Throat Pain
- Cold & Flu

Reduces fever.

Warnings
Do not use this product:
- If you are allergic to ibuprofen, aspirin or other anti-inflammatory medicine, pain reliever / fever reducer or any other component of this product
- If you are pregnant, or trying to become pregnant
- If you have a current or past history of stomach ulcer / bleeding / disorders
- If you have impaired kidney function or heart failure
- In children under 12 years

Unless your doctor has told you do not use this product:
- If you have asthma, high blood pressure, heart or liver disease
- If you are breastfeeding
- With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
- If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin
- If you are 65 years or over or under a doctor's care for any serious condition

While using this product:
- Stop use and ask your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
- Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
- Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage
- Side effects may be minimised by using the smallest dose for the shortest duration of time
- In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)

Contains sulfites, soya bean products, and potassium (17.84 mg per capsule)

Directions For Use
Adults and children 12 years of age and over.
Take 1 or 2 capsules, swallowed whole with water.
Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 capsules in 24 hours.

Continued...

VX

20 SOFT CAPSULES

RELIEVES PAIN FAST

Advil LIQUID CAPS

HALEON

RELIEVES PAIN FAST

20 SOFT CAPSULES

AUST R 471310

Advil LIQUID CAPS

Ibuprofen 200 mg

HEADACHE • PERIOD PAIN
MUSCLE PAIN • BACK PAIN

9 310488 017157

Medicine Information (continued)
Other Information:
STORE BELOW 30°C
IN A DRY PLACE

Do not use this product if the foil is broken

Haleon
Sydney, NSW, Australia & Auckland, New Zealand
AU: 1800 028 533
NZ: 0800 540 144
www.advil.net.au

Trade Marks owned or licensed by Haleon.
©2024 Haleon or licensor.

KEEP OUT OF REACH OF CHILDREN

B. No.:

EXP. :

02153944