Medicine Information (continued)

Unless your doctor has told you do not use this product:

- If you are 65 years or over or under a doctor's care for any serious condition

While using this product:

- Stop use and see your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
- Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage
- Side effects may be minimised by using the smallest dose for the shortest duration of time
- Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
- In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)

Contains sulfites, soya bean products, galactose and 15.9 mg of potassium per capsule

Directions For Use

For use by Adults and Children from 12 years and over. Take 1 or 2 capsules swallowed whole with water. Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 capsules in 24 hours.

Continued on panel below...

Each Advil Minis Liquid Capsule contains liquid ibuprofen for fast pain relief

Medicine Information (continued)

Other Information:
STORE BELOW 25°C
Do not use if foil seal is broken or missing

Haleon
Sydney, NSW,
Australia &
Auckland,
New Zealand

AU: 1800 028 533
NZ: 0800 540 144
www.advil.net.au

KEEP OUT OF REACH OF CHILDREN

Trade Marks owned or licensed by Haleon.
©2024 Haleon or licensor.

62000000206843

EXP.

62000000206843

HALEON

Advil

Ibuprofen 200 mg

minis

Relieves Pain FAST

20 Liquid Capsules

AUST R 311280

Advil minis

9 310488 004263

Medicine Information

Active Ingredient (in each liquid capsule)
Ibuprofen 200 mg

Uses

Advil provides temporary relief of pain, discomfort and/or inflammation associated with:

- Headache
- Migraine Headache
- Tension Headache
- Period Pain
- Sore Throat Pain
- Back Pain
- Dental Pain
- Cold & Flu Symptoms
- Muscle Pain
- Arthritis Pain

Advil reduces fever

Warnings

Do not use this product:

- If you are allergic to ibuprofen, aspirin or other anti-inflammatory medicine, pain reliever / fever reducer or any other component of this product
- If you are pregnant, or trying to become pregnant
- If you have a current or past history of stomach ulcer / bleeding / disorders
- If you have impaired kidney function or heart failure
- In children under 12 years

Unless your doctor has told you do not use this product:

- If you have asthma, high blood pressure, heart or liver disease
- If you are breast feeding
- With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
- If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin

Continued on panel below...

Medicine Information (continued)

Unless your doctor has told you do not use this product:

- If you are 65 years or over or under a doctor's care for any serious condition

While using this product:

- Stop use and see your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
- Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage
- Side effects may be minimised by using the smallest dose for the shortest duration of time
- Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
- In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)

Contains sulfites, soya bean products, galactose and 15.9 mg of potassium per capsule

Directions For Use

For use by Adults and Children from 12 years and over. Take 1 or 2 capsules swallowed whole with water. Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 capsules in 24 hours.

Continued on panel below...

Bottle Inside

Each Advil Minis Soft Capsule contains liquid ibuprofen for fast pain relief

Medicine Information (continued)

Other Information:
STORE BELOW 25°C
Do not use if foil seal is broken or missing

Haleon
Sydney, NSW, Australia
& Auckland, New Zealand

AU: 1800 028 533
NZ: 0800 540 144
www.advil.net.au

KEEP OUT OF REACH OF CHILDREN

Trade Marks owned or licensed by Haleon.
©2024 Haleon or licensor.

NPA016450

Advil

HALEON

Advil

Ibuprofen 200mg

minis

RELIEVES PAIN FAST

AUST R 311280

20 SOFT CAPSULES

9 310488 004263

Ⓑ

EXP.

NPA016450

Medicine Information

Active Ingredient (in each soft capsule)
Ibuprofen 200 mg

Uses

Advil provides temporary relief of pain, discomfort and/or inflammation associated with:

- Headache
- Migraine Headache
- Tension Headache
- Period Pain
- Sore Throat Pain
- Back Pain
- Dental Pain
- Cold & Flu Symptoms
- Muscle Pain
- Arthritis Pain

Advil reduces fever

Warnings

Do not use this product:

- If you are allergic to ibuprofen, aspirin or other anti-inflammatory medicine, pain reliever / fever reducer or any other component of this product
- If you are pregnant, or trying to become pregnant
- If you have a current or past history of stomach ulcer / bleeding / disorders
- If you have impaired kidney function or heart failure
- In children under 12 years

Unless your doctor has told you do not use this product:

- If you have asthma, high blood pressure, heart or liver disease
- If you are breast feeding
- With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
- If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin

Continued on panel below...