LOT:
EXP:
HALEON
Advil
Relieves Pain FAST
24 Tablets
EASY TO SWALLOW
Advil
Ibuprofen 200 mg
Relieves Pain FAST
• HEADACHE • BACK PAIN
• MUSCLE PAIN • PERIOD PAIN
AUST R 104221
9 310488 017065
Made in Italy
Medicine Information (continued)
Other Information
STORE BELOW 25°C
KEEP OUT OF REACH OF CHILDREN
Haleon
Sydney, NSW, Australia & Auckland, New Zealand
AU: 1800 028 533
NZ: 0800 540 144
www.advil.net.au
Trade Marks owned or licensed by Haleon.
©2024 Haleon or licensor.
Advil
Relieves Pain FAST
62000000208509
Medicine Information
Active Ingredient (in each tablet)
Ibuprofen 200 mg
Uses
Advil provides temporary relief of pain and discomfort from:
• Headache • Period Pain • Sore Throat Pain
• Back Pain • Dental Pain • Cold & Flu
• Muscle Pain • Arthritis Pain
Advil reduces fever.
Warnings
Do not use this product:
• If you are allergic to ibuprofen, aspirin or other anti-inflammatory medicine, pain reliever / fever reducer or any other component of this product
• If you are pregnant, or trying to become pregnant
• If you have a current or past history of stomach ulcer / bleeding / disorders
• If you have impaired kidney function or heart failure
• If the foil seal is broken
Unless your doctor has told you do not use this product:
• If you have asthma, high blood pressure, heart or liver disease
• If you are breast feeding
• With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
• If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin
• If you are 65 years or over or under a doctor's care for any serious condition
While using this product:
• Stop use and ask your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
• Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
• Do not use for more than a few days at a time unless a doctor has told you to. Do not exceed the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage
• Side effects may be minimised by using the smallest dose for the shortest duration of time
• In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)
Contains sugars, benzoates, hydroxybenzoates, sulfites, and soya beans or soya bean products
Directions For Use
For use by Adults and Children from 12 years and over.
Begin by taking 1 or 2 tablets, swallowed whole with water. One tablet will often be adequate for mild pain, but for more severe pain you may require two tablets.
Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 tablets in 24 hours. Do not give to children under 12 years.
Continued...

XXXXXX

XXXXXX