Superseded Artwork



New Artwork

XXXXXX

EXP: :101 24 TABLETS RELIEVES PAIN **FAST** Hateon
Sydney, NSW, Australia & Auckland, Ne
L AU: 1800 028 533
NZ: 0800 540 144

Www.advil.net_au
Tade Marks owned or licensed by Hates RELIEVES PAIN FAST

• HEADACHE
• PERIOD PAIN
• MUSCLE PAIN • BACK PAIN Ibuprofen 200mg Medicine Information (continued) Other Information STORE BELOW 25°C KEEP OUT OF REACH OF CHILDREN NPA017019

Medicine Information

Active Ingredient (in each tablet)

Uses Advil provides temporary relief of pain and discomfort from:

- Headache
 Back Pain
 Muscle Pain Period Pain
 Dental Pain
 Arthritis Pain
 - Sore Throat Pain
 Cold & Flu

- Warnings
 Do not use this product:

 If you are altergic to bluprofen, aspirin or other anti-inflammatory medicine, pain reliever? fever reducer or any other component of this product

 If you are pregnant, or trying to become pregnant

 If you have a current or past history of stomach ulcer / bleeding / disorders

 If you have impaired kidney function or heart failure

- If the foil seal is broken

- Unless your doctor has told you do not use this product:

 If you have asthma, high blood pressure, heart or liver disease

 If you are breast feeding

- If you are breast feeding
 With other medicines containing ibuprofen, aspirin, other anti-inflammatory medicines, blood thinners or any other medicines you take regularly
 If you are taking aspirin for heart attack or stroke as ibuprofen may decrease this benefit of aspirin
 If you are 65 years or over or under a doctor's care for any serious condition
 White using this product:
 Stop use and ask your doctor immediately if you see an allergic reaction including skin reddening, rash or blisters or if pain or fever gets worse or lasts for more than a few days
 Stop use and ask the doctor if you see blood in your vomit or have bloody or black stools
 On on tuse for more than a few days at a time unless a doctor has told you to.
 Do not exceed the recommended dose. Excessive or prolonged use can be harmful.
- To mote see the recommended dose. Excessive or prolonged use can be harmful and increase the risk of heart attack, heart disease, stroke or liver damage. Side effects hay be minimised by using the smallest dose for the shortest duration of time.
- In case of accidental overdose, stop use and contact immediately the Poisons Information Centre (Australia 13 11 26, New Zealand 0800 764 766)
 Contains sugars, benzoates, hydroxybenzoates, sulfites, and soya beans or soya bean products

Directions For Use
For use by Adults and Children from 12 years and over.
Begin by taking 1 or 2 tablets, swallowed whole with water. One tablet will
often be adequate for mild pain, but for more severe pain you may require

Repeat dose if necessary after 4 to 6 hours. Do not exceed 6 tablets in 24 hours. Do not give to children under 12 years.

Continued...

XXXXXX