



For Children

Fever, Symptoms of Cold and Pain

For Relief of fever & pain.

Indications

Panadol For Children is indicated for the relief of fever, headache, symptoms of cold and flu, teething pain and toothache.

Description

Each tablet contains Paracetamol 120mg, chewable and cherry flavoured.

Excipients: Cherry Flavour Powder, D&C Red Lake Blend, Mannitol, Saccharin Sodium, Starch, Stearic Acid, Ethylcellulose.

Dosage and administration

For oral administration only.

Children aged 2 years and above:

Age	Ave. Weight	Dose
2-4 yrs	12-16kg	1.5 tabs
4-7 yrs	16-21.5kg	2 tabs
7-9 yrs	21.5-25kg	2.5 tabs
9-10 yrs	25-28.7kg	3 tabs
10-11 yrs	28.7-33kg	3.5 tabs
11-12 yrs	33-37.5kg	4 tabs

Do not exceed the stated dose. Do not use with this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu. Do not take more than four doses in any 24 hour period. Do not take more frequently than once every 4 hours. Do not take for more than 3 days without asking your doctor. Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary. Not recommended in children under 2 years.

Tablets may be chewed, or dissolved in water or fruit juice. For children below 6 years, tablets should be mixed with water or fruit juice.

Contraindications

Do not use if: The child is allergic to Paracetamol or any of the other ingredients in the product.

Warnings and Precautions

Contains Paracetamol. Do not take more than the recommended dose as it may cause serious harm to your child's liver. Do not use this medicine if your child is taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep. Always read and follow the label.

Check with the doctor before use if:

- The child has liver or kidney problems.
- The child is underweight or malnourished.

HALEON

[NPA018199]



Your child may need to avoid using this product altogether or limit the amount of paracetamol that your child takes.

The child has a severe infection, is severely malnourished, severely underweight as this may increase the risk of metabolic acidosis.

- Signs of metabolic acidosis:
 - deep, rapid, difficult breathing
 - feeling sick (nausea)
 - being sick (vomiting)
 - loss of appetite

Contact your doctor immediately if the child gets a combination of these symptoms.

- The child had a serious skin reaction with this medicine, do not take it or any products containing paracetamol or acetaminophen again. Doing so could cause the child to have another serious skin reaction.

Please see the doctor if the child's symptoms do not improve.

Keep out of reach and sight of children.

Drug Interaction

Before taking this medication, make sure you consult the doctor if the child is taking Warfarin or similar medicines used to thin blood. Chronic consumption of anticonvulsants may reduce the efficiency of paracetamol by increasing its clearance through the liver.

Adverse Reactions

Stop taking this medicine and tell the doctor immediately if:

- The child experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- The child experience any skin rash or reaction. Symptoms of serious skin reactions may include skin reddening, rash, blisters, and separation of the upper surface of the skin from the lower layers. These serious skin reactions are rare, but can occur even if you have taken this medicine in the past without any problems.
- The child has previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experiences a similar reaction with this product.
- The child experiences unexplained bruising or bleeding.
- The child is becoming unusually tired and getting more infections (such as colds) than usual.
- The child experience mouth ulcers.

These reactions are rare.

Overdose

If the child takes more of the medicine than he/she should: Seek medical advice immediately even if the child do not have any symptoms because of the risk of liver failure.

Pack Size

Box of 24's

Storage Conditions

Store below 30°C.

For further information, please refer to your doctor or pharmacist.

GDS Version : 7.0

Last updated : October 2024.

Trade Marks are owned by or licensed to the Haleon group of companies.

©2025 Haleon group of companies or its licensor.

[NPA018199]

