



Panadol CR
120mg/5ml

WHAT IS PANADOL CHILDREN SUSPENSION USED FOR?

Panadol Children Suspension is indicated for the relief of fever and pain associated with headache, teething, toothache, earache, immunization, symptoms of cold and flu.

WHAT DOES PANADOL CHILDREN SUSPENSION CONTAIN?

Each teaspoon (5ml) contains Paracetamol 120mg.

Excipients: Sucrose Crystalline, Sorbitol Solution 70%, Glycerin, Xanthan Gum, Strawberry Flavour, Methyl Hydroxybenzoate, Carmoisine, Purified water.

APPEARANCE

A pink suspension with strawberry odour.

DOSAGE AND ADMINISTRATION

- For oral administration only.
- This product is intended for children age 1 month and above. For children weighing above 17kg, use Panadol Children Suspension 250mg/5ml.
- Shake well before use.

Recommended dosage of paracetamol per dose is 10-15mg/kg of body weight.

WEIGHT (kg)	DOSE (ml)
4 – 5 kg	2.5 ml
6 – 7 kg	3.5 ml
8 – 9 kg	5 ml
10 – 11 kg	6 ml
12 – 13 kg	7.5 ml
14 – 15 kg	8.5 ml
16 – 17 kg	10 ml

- Do not exceed the stated dose.
- Use the smallest dose that you need to treat your child's symptoms and use the medicine for the shortest period of time necessary.
- Do not use this medicine if your child is taking any other prescription or non-prescription medicine containing paracetamol to treat pain, fever, symptoms of cold and flu.
- Do not take more than four doses in any 24 hour period.
- Do not take more frequently than every 4 hours.
- Do not take more than 3 days without asking your doctor.
- Not recommended for children under 1 month.
- For children below 4 kg: seek medical advice before the use of the product for appropriate dose and administration.

CONTRAINDICATIONS

Do not use if: The child is allergic to Paracetamol or any of the other ingredients in the product.

WARNINGS AND PRECAUTIONS

Contains Paracetamol. Do not take more than the recommended dose as it may cause serious harm to your child's liver. Always read and follow the label.

Check with the doctor before use if the child:

- has liver or kidney problems.
- is underweight or malnourished.

The child may need to avoid using this product altogether or limit the amount of paracetamol that the child take.

- has a severe infection, is severely malnourished, severely underweight as this may increase the risk of metabolic acidosis.

Signs of metabolic acidosis include:

→ deep, rapid, difficult breathing

→ feeling sick (nausea), being sick (vomiting)

→ loss of appetite

Contact a doctor immediately if your child gets a combination of these symptoms.

- If the child have had a serious skin reaction with this medicine, do not take it or any products containing paracetamol or acetaminophen again. Doing so could cause the child to have another serious skin reaction.

Please see the doctor if the child's symptoms do not improve.

Keep out of sight and reach of children.

This product contains Carmoisine which may cause allergic-type reactions. Patients

with rare hereditary problems of fructose intolerance should not take this medicine.

This medicine may have a mild laxative effect and cause gastrointestinal discomfort. This medicine contains 150 mg oxibutol in each ml. If you have been told by your

This medicine contains 450.30mg sorbitol in each ml. If you have been told by your doctor that your child has an intolerance to some sugars, contact your doctor.

doctor that your child has an intolerance to some sugars, contact your doctor before taking this medicinal product.

DRUG INTERACTIONS

Before taking this medicine, make sure you consult the doctor if the child is taking Warfarin or similar medicines used to thin the blood. Chronic consumption of anticonvulsants may reduce the efficiency of paracetamol by increasing its clearance through the liver.

OVERDOSE

If the child takes more of the medicine than he/she should: Seek medical advice immediately even if the child does not have any symptoms because of the risk of liver failure.

HALEON

- ADVERSE REACTIONS**
Stop taking this medicine and tell the doctor immediately if:
- The child experiences allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
 - The child experiences any skin rash or reaction. Symptoms of serious skin reactions may include skin reddening, rash, blisters, and separation of the upper surface of the skin from the lower layers. These skin reactions are rare, but can occur even if you have taken this medicine in the past without any problems.
 - The child has previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experiences a similar reaction with this product.
 - The child experience unexplained bruising or bleeding.
 - The child is becoming unusually tired and getting more infections (such as colds) than usual.
 - The child experience mouth ulcers.

These reactions are rare.

STORAGE CONDITION

- Store at or below 30°C
- Shelf life after first opening: 12 months
- Protect from light

Do not use if cap is broken or separated.
For further information, ask a doctor or pharmacist.
GDS Version: 9.0
Last updated: April 2024

Panadol Children Suspension 120mg/5ml is available in Child Safety Caps.



Trademarks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies.

HALEON
NPA214887