




• You have high blood pressure, heart disease or blood vessel disease such as Raynaud's phenomenon (which may appear as pain, numbness, tingling and color change (white, blue then red) pain in the fingers or toes when exposed to cold), diabetes (too much sugar in your blood) or hyperthyroidism (an overactive thyroid gland). • You have glaucoma (excessive pressure inside your eyes). • You have phaeochromocytoma (a rare tumour of the adrenal glands which sit above the kidneys). • You have difficulty urinating due to an enlarged prostate gland. • If you have had a serious skin reaction with this medicine, do not take it or any products containing paracetamol or acetaminophen again. Doing so could cause you to have another serious skin reaction. • You are pregnant, trying to become pregnant or breastfeeding. Before taking this medicine, make sure you consult your doctor if you are taking any of the following: • Warfarin or similar medicines used to thin the blood. • Medicines to control your blood pressure, such as beta-blockers. • Digoxin or similar medicines for heart disease. • Appetite suppressants or stimulants. • Medicines to treat depression such as tricyclic antidepressants eg. Amitriptyline. • Medicines for the treatment of acute migraine headache (e.g. ergotamine, methysergide). You should also tell your doctor or pharmacist about any other medicines you are taking. If your symptoms worsen, do not improve within 7 days, or are accompanied by high fever, rash or persistent headache, consult your doctor. Keep out of sight and reach of children. Overdose: If you take more of the medicine than you should: Seek medical advice immediately even if you feel well because of the risk of liver failure. Ability to perform tasks that require judgement, motor or cognitive skills: This product may cause dizziness. If affected, do not drive or operate machinery. Side Effects: When using this product, you may experience: headache, dizziness, nervousness, difficulty sleeping, increased blood pressure, nausea and vomiting. Stop taking this medicine and tell your doctor immediately if: • You experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face. • You experience any skin rash or reaction. Symptoms of serious skin reactions may include skin reddening, rash, blisters, and separation of the upper surface of the skin from the lower layers. These serious skin reactions are rare, but can occur even if you have taken this medicine in the past without any problems. • You have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product. • You experience unexplained bruising or bleeding. • You experience an unusually fast pulse rate, or a sensation of an unusually fast or irregular heartbeat. • You experience loss of vision, which may be due to abnormally high blood pressure in the eye. This is more likely to occur in those with glaucoma. • You experience difficulty in passing water. This is more likely to occur in men with an enlarged prostate gland. • You experience mouth ulcers. These reactions are rare or very rare. Storage conditions: Store below 30°C, away from direct sunlight and in a cool place. Use only if foil seal is intact. For further information, ask a doctor or pharmacist. GDS Version: 7.0 Date Updated : October 2024 Trade Marks are owned by or licensed to the Haleon group of companies. ©2024 Haleon group of companies or its licensor.	SIN13878P 9001028
---	------------------------------