	harma code Ref. No. 3217	1	
		교	
Pana	dol Extend gsk	PEEL HERE TO REVEAL PRODUCT INFORMATION	
INDICATI			
associate	ixtend is used to give long lasting relief of pain d with osteoarthritis, backaches, mild to moderate joint	ION CONTRACT	
pains and injuries.	muscular aches such as sprains, strains, and sports	-	
Panadol E	Extend is also suitable for headache and migraine.		
	Each sustained release caplet contains 665mg of Paracetamol. The caplet also contains Hypromellose,		
	Pregelatinised Starch, Povidone, Magnesium Stearate, Croscarmellose Sodium, Stearic Acid,		
	Triacetin, Carnauba Wax. Caplet is a white to off-white capsule shaped		
	film-coated tablet debossed with "8" on one side and blank on the other.		
	DOSAGE		
	For oral administration only. Do not exceed the stated dose. Use the smallest dose that you need to		
	treat your symptoms and use the medicine for the shortest period of time necessary. Do not take more		
are taking	y than every 6 hours. Do not use this medicine if you any other prescription or non-prescription medicines		
and flu.	g paracetamol to treat pain, fever, symptoms of cold		
Take 2 ca	cluding the elderly) and children over 12 years: plets three times a day, every 6-8 hours as required.		
Children u	ke more than 6 caplets per day. Inder 12 years: Not recommended for children under		
-	f 12 years. EREACTIONS		
	ng this medicine and tell your doctor immediately if: operience allergic reactions such as skin rash or		
), sometimes with breathing problems or swelling of s, tongue, throat or face.		
seriou	perience any skin rash or reaction. Symptoms of s skin reactions may include skin reddening, rash,		
	s and separation of the upper surface of the skin from ver layers. These serious skin reactions are rare, but		
	cur even if you have taken this medicine in the past it any problems.		
	xperience mouth ulcers. ave previously experienced breathing problems with		
	or non-steroidal anti-inflammatories, and experience lar reaction with this product.		
	perience unexplained bruising or bleeding. Ictions are rare.		
PREGNAM Talk to yo	ICY ur health care professional before taking paracetamol		
	pregnant. taking the lowest effective dose for the shortest period		
	ou can take this product whilst breastfeeding.		
Before tal	king this medicine, make sure you consult your doctor taking Warfarin or similar medicines used to thin the		
blood.	,		
Do not us	NDICATIONS e if you are allergic to Paracetamol or any of the other is the generation		
ingredien	ts in the product. SS AND PRECAUTIONS		
WARNING	o much paracetamol can cause serious harm to		
Taking to	Always read and follow the label.		
Taking to your liver. Check wit	Always read and follow the label. th your doctor before use if: ave liver or kidney problems.		
Taking to your liver Check wii • You ha • You ar • You ar	th you ^r doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol.		
Taking to your liver. Check wi • You ha • You or You re You may limit the a	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or unount of paracetamol that you take. If you have		
Taking to your liver Check wii • You ha • You re You rey limit the a a severe i underwei	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or imount of paracetamol that you take. If you have nfection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may		
Taking to your liver. Check wii • You ar • You re You rey Jumit the a a severe i underwei increase f	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or mount of paracetamol that you take. If you have nfection, are severely malnourished, severely		
Taking tor your liver. Check wii • You ha • You ra • You ra You ray limit the a a severe i underwei increase f • Signs • Deep f	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or imount of paracetamol that you take. If you have infection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis.		
Taking to your liver. Check wil • You ha • You re You rey You may limit the a a severe i underwei increase t • Signs - Deep - Feelin - Feelin	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or unount of paracetamol that you take. If you have infection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis. of metabolic acidosis include: rapid, difficult breathing		
Taking to your liver. Check wii • You ar • You re You may limit the a a severe i underwei increase • Signs - Deep - Feelin - Loss o Contact a	th you ^r doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or imount of paracetamol that you take. If you have nfection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis. of metabolic acidosis include: rapid, difficult breathing g sick (nausea), being sick (vomiting) f appetite		
Taking to your liver. Check wil • You ha • You re You re You rey limit the a a severe i underwei increase f • Signs - Deep - Feelin - Loss o Contact a these syn improve. If you hav	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or umount of paracetamol that you take. If you have nfection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis. of metabolic acidosis include: rapid, difficult breathing g sick (nausea), being sick (vomiting) f appetite doctor immediately if you get a combination of		
Taking tor your liver, Check wii • You ar • You re You may limit the a a severe i underwei increase f • Signs - Deep - Feelin - Loss o Contact a these syn improve. If you hav do not tal acetamin	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or imount of paracetamol that you take. If you have nfection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis. of metabolic acidosis include: rapid, difficult breathing g sick (nausea), being sick (vomiting) f appetite doctor immediately if you get a combination of nptoms. Please see a doctor if your symptoms do not e had a serious skin reaction with this medicine,		
Taking to your liver. Check wil • You ha • You re You may limit the a a severe i underwei increase 1 • Signs • Deep - Feelin • Loss o Contact a these syn improve. If you hav do not tal acetamin serious sl Please se	th your doctor before use if: ave liver or kidney problems. e severely underweight or malnourished. gularly drink alcohol. need to avoid using this product altogether or umount of paracetamol that you take. If you have nfection, are severely malnourished, severely ght, or are a chronic heavy alcohol user , this may the risk of metabolic acidosis. of metabolic acidosis include: rapid, difficult breathing g sick (nausea), being sick (vomiting) f appetite doctor immediately if you get a combination of nptoms. Please see a doctor if your symptoms do not e had a serious skin reaction with this medicine, ke it or any products containing paracetamol or opphen again. Doing so could cause you to have another		


Item Number: P3217P (used with Manufacturing Sin GSK-AUS-Ermingt Market or Pack Or CH-Singapore-SGI Market Trade Nam Panadol Extend Colour Standard F Number: N/A Technical Referen ELEAF0012 (273.0 (do NOT include the technical ref Printing Process: Flexography Substrate: White Paper	te: ton-Al wner: P ne: Refere ce No 5 x 83	JMCA nce (s).: mm)
GSK-AUS-Ermingt Market or Pack Or CH-Singapore-SG Market Trade Nan Panadol Extend Colour Standard F Number: N/A Technical Referen ELEAF0012 (273.0 (do NOT include the technical ref Printing Process: Flexography Substrate:	ton-Al wner: P ne: Refere ce Not	nce (s).: mm)
CH-Singapore-SG Market Trade Nan Panadol Extend Colour Standard F Number: N/A Technical Referen ELEAF0012 (273.0 (do NOT include the technical ref Printing Process: Flexography Substrate:	P ne: Refere ce No 5 x 83	(s).: mm)
Panadol Extend Colour Standard F Number: N/A Technical Referen ELEAF0012 (273.0 (do NOT include the technical ref Printing Process: Flexography Substrate:	Refere ce No 5 x 83	(s).: mm)
Number: N/A Technical Referen ELEAF0012 (273.0 ^(do NOT include the technical reference) Printing Process: Flexography Substrate:	ce No 5 x 83	(s).: mm)
ELEAF0012 (273.0 (do NOT include the technical ref Printing Process: Flexography Substrate:	5 x 83	mm)
Flexography Substrate:		
Colours	Fotal: 1	-
К		
Varnishes 1	Fotal: 0	
Special Finishes	Fotal: 0)
Artwork copyright is t	•	
GlaxoSmithKline Gro All suppliers providing a ser components of any descrip they have a licence for all f conjunction with The distribution and use of f a licence constitutes an i infringement. GSK will not ar	pup of Co rvice to GS otion must fonts / soft GSK artwo fonts / sof intellectua	mpanies 5K for printed : ensure that ware used in ork. tware without Il property
breach of third party intelle printed compone The GSK certification / audit p to declare that they do not software and may require t evidence of such li	ctual prop ent supplie process ree t use unlic the supplie	perty rights by ers. quires suppliers ensed fonts / er to produce
ATTENTION • To Ensure Accurate PDF V		ENTION
Io ensure Accurate PDV FOR SCREEN VIEWING: Professional or Adobe Acrol Professional (higher than 7 must be activated for accur FOR PRINTING: Use only version 7 or higher. "Apply "Simulate Overprinting" print settings for printing	Use Adob bat Reade). Overpr rate on scr Acrobat F Overprin must be a	e Acrobat 7 r, Standard or int Preview een viewing. Professional t Preview " or ctivated in the

GSK Market is responsible
for this product, its design
and content.TEXT SIZE CONTAINED IN THIS ARTWORKEnsure the artwork is
thoroughly checked, all the
text proof-read and approveLeading: 8.0ptRSC GSK is responsible for si
technical requirements and
pre-press suitability.

Microtext: N

Ermington – Additional Artwork information Panel		
Pharmacode (or 2D Matrix) Number	3217 (Pharmacode)	
Varnish Type	N/A	
Specification number	ESPLEAF0011	

for this product, its design and content. Ensure the artwork is thoroughly checked, all the text proof-read and approved. RSC GSK is responsible for site technical requirements and pre-press suitability. GSK Market is responsible to advise RSC in case changes required impact the followings: Formulation Tablet embossing Storage conditions Shelf Life



GlaxoSmithKlir Artwork Inforn Panel		RSC A/W Version: 1	
Item Number: P3217P (used w	ith C321	16P)	_
Manufacturing GSK-AUS-Ermin		JMCA	
Market or Pack CH-Singapore-S			
Market Trade Na Panadol Extend			
Colour Standarc Number: N/A	l Refere	nce	1
Technical Refere	.05 x 83	mm)	
Printing Process Flexography	:		
Substrate: White Paper			g Bar
Colours	Total: 1	•	180 mm Measuring Bai
к			n Mea
			80 mn
Varnishes	Total: 0		₩
			ion 2
			NOV_2013 Version 2
Special Finishes	Total: 0)	1
			AIP_Disclaimers_INDD
			AIP_Disc
Artwork copyright i GlaxoSmithKline C All suppliers providing a components of any dess they have a licence for a conjunction wi The distribution and use e a licence constitutes a infringement. GSK will no breach of third party inte printed compo The GSK certification / aud to declare that they do software and may require evidence of suc	Froup of Con service to GS cription must lil fonts / soft th GSK artwo of fonts / soft nn intellectual t accept any ellectual proponent supplie lit process rec not use unlic.	mpanies K for printed ensure that ware used in ork. I property liability for the erty rights by ers. quires suppliers ensed fonts / er to produce	-
		ENTION	
To Ensure Accurate PD FOR SCREEN VIEWINN Professional or Adobe Ac Professional (higher tha must be activated for ac FOR PRINTING: Use on version 7 or higher. "App "Simulate Overprinting print settings for printi	G: Use Adobe crobat Reade n 7). Overpr curate on scr nly Acrobat F ly Overprin g" must be a	e Acrobat 7 r, Standard or int Preview een viewing. Professional t Preview " or ctivated in the	
	RT/	NT	

for this product, its design and content. Ensure the artwork is thoroughly checked, all the text proof-read and approved.

RSC GSK is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise RSC in case changes required impact the followings: Formulation Tablet embossing Storage conditions Shelf Life

TEXT SIZE CONTAINED IN THIS ARTWORK Body text size: 7.0pt Leading: 8.0pt Horizontal Scale: 100% Smallest text size: 7.0pt Microtext: N

Ermington – Additional Artwork information Panel		
Pharmacode (or 2D Matrix) Number	3217 (Pharmacode)	
Varnish Type	N/A	
Specification number	ESPLEAF0011	