



e-Banner	Project: CO-0002355	Document: PPC-0001277	Version: 2
	Site Code: 911719	Operator: WD809459	Date/Time Created: 23.Jan.2018 08:57 GMT





Panadol For Children

Fever, Symptoms of Cold and Pain

For Relief of fever & pain.

Indications
Panadol For Children is indicated for the relief of fever, headache, symptoms of cold and flu, teething pain and toothache.

Description
Each tablet contains Paracetamol 120mg, chewable and cherry flavoured.
Excipients: Cherry Flavour Powder, Erythrosine, Mannitol, Saccharin Sodium, Starch, Stearic Acid, Ethylcellulose.

Dosage and administration
Oral administration only.
Children aged 2 years and above:

Age	Ave. Weight	Dose
2-4 yrs	12-16kg	1.5 tabs
4-7 yrs	16-21.5kg	2 tabs
7-9 yrs	21.5-25kg	2.5 tabs
9-10 yrs	25-28.7kg	3 tabs
10-11 yrs	28.7-33kg	3.5 tabs
11-12 yrs	33-37.5kg	4 tabs

Do not exceed the stated dose. Do not use with this medicine if you are taking any other prescription or non-prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu. Do not take more than four doses in any 24 hour period. Do not take more frequently than once every 4 hours. Do not take for more than 3 days without asking your doctor. Use the smallest dose that you need to treat your symptoms and use the medicine for the shortest period of time necessary. Not recommended in children under 2 years.
Tablets may be chewed, or dissolved in water or fruit juice. For children below 6 years, tablets should be mixed with water or fruit juice.


Pack Size
Box of 24's and 144's.

Expiry
Do not use beyond the expiry date.

Adverse Reactions
Stop taking this medicine and tell the doctor immediately if:

- The child experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- The child experience any skin rash or reaction. Symptoms of serious skin reactions may





include skin reddening, rash, blisters, and separation of the upper surface of the skin from the lower layers. These serious skin reactions are rare, but can occur even if you have taken this medicine in the past without any problems.

- The child has previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experiences a similar reaction with this product.
- The child experiences unexplained bruising or bleeding.
- The child is becoming unusually tired and getting more infections (such as colds) than usual.
- The child experience mouth ulcers.

These reactions are rare.

Contraindications
Do not use if: The child is allergic to Paracetamol or any of the other ingredients in the product.

Warnings and Precautions
Taking too much paracetamol can cause serious harm to your liver. Always read and follow the label. Check with the doctor before use if:

- The child has liver or kidney problems.
- The child is underweight or malnourished.

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.
The child has a severe infection, is severely malnourished, severely underweight as this may increase the risk of metabolic acidosis.

Signs of metabolic acidosis:

- deep, rapid difficult breathing
- feeling sick (nausea)
- being sick (vomiting)
- loss of appetite

Contact your doctor immediately if the child gets a combination of these symptoms.
The child had a serious skin reaction with this medicine, do not take it or any products containing paracetamol or acetaminophen again. Doing so could cause the child to have another serious skin reaction.

Please see the doctor if the child's symptoms do not improve.
Keep out of reach and sight of children.

PREGNANCY
Talk to your health care professional before taking paracetamol if you are pregnant.
Consider taking the lowest effective dose for the shortest period of time. You can take this product whilst breastfeeding.

Drug Interaction
Before taking this medication, make sure you consult the doctor if the child is taking Warfarin or similar medicines used to thin blood. Chronic consumption of anticonvulsants may reduce the efficiency of paracetamol by increasing its clearance through the liver.

Overdose
If the child takes more of the medicine than he/she should: Seek medical advice immediately even if the child do not have any symptoms because of the risk of liver failure.

Storage Conditions
Store below 30°C.
For further information, please refer to your doctor or pharmacist.
Trade marks are owned by or licensed to the GSK group of companies.
GDS Version : 7.0
Last updated : October 2017
© 2016 GSK group of companies. All Rights Reserved.

Version: 1		Harmony AMS - Artwork Information Panel		gsk	
Manufacturing Site Number: 911719	Manufacturing Site(s): GSK_KUALA LUMPUR_MALAYSIA	Technical Drawing (Do NOT include version number): IMA_INS_120X180_DRW			
Product Market Trade Name: Panadol Children	Approving Market(s): Singapore-SGP	Material Spec. (Do NOT include version number): MSP-1500032			
Print Process: Offset	Colour Standard Reference: PLEASE REFER TO INDUSTRY STANDARD PANTONE VALUES	Material Type: Paper/Cardboard Leaflet paper	SUBSTRATE		
Total Colours & Varnishes: 1		Body Text Size: 7.0pt	Smallest Text Size: 7.0pt		
BLACK		Leading: 8.0pt			
Total Special Finishes: 0		Horizontal Scale: 100%			
		Microtext: N			
Additional Info (1): N/A		Additional Info (2): N/A		Additional Info (3): N/A	
<small>Artwork copyright is the property of the GSK Group of Companies. All suppliers providing a service to GSK for printed components of any description must ensure that they have a licence for all fonts / software used in conjunction with GSK artwork. The distribution and use of fonts / software without a licence constitutes an intellectual property infringement. GSK will not accept any liability for the breach of third party intellectual property rights by printed component suppliers. The GSK certification / audit process requires suppliers to declare that they do not use unlicensed fonts / software and may require the supplier to produce evidence of such licence to GSK.</small>					
200 mm Measuring Bar					
If an e-banner DOES NOT appear on the top of this document, THEN this document has NOT been printed from the Harmony system.					

IMPORTANT

GSK LOC is responsible to approve the change documentation, artwork brief and final artwork, ensuring that it is accurate, consistent and complete.

GSK SDC is responsible for site technical requirements and pre-press suitability.

GSK Market is responsible to advise SDC when changes required impact the following:

Formulation
Tablet embossing
Storage conditions
Shelf Life

NOTE TO MARKET

Local approvers must ensure that trade mark and copyright statements included in the brief comply with guidance provided by Legal: Global Trade Marks.