

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**  
Ibuprofen 200 mg (NSAID)\* ..... Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.  
Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

For most recent product information, visit [www.Advii.com](http://www.Advii.com). ©2021 GSK or licensor. Trademarks owned or licensed by GSK.

**Drug Facts (continued)**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** **if** you have ever had an allergic reaction to any other pain reliever/fever reducer **or** right before or after heart surgery

**Ask a doctor before use if** **if** you have problems or serious side effects from taking pain relievers or fever reducers **or** you have a history of stomach problems, such as heartburn **or** you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke **or** you are taking a diuretic

**Ask a doctor or pharmacist before use if you are** **if** you are under a doctor's care for any serious condition **or** taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin **or** taking any other drug

**When using this product**  
take with food or milk if stomach upset occurs

128 CODE AREA

128 CODE AREA

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**if pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

128 CODE AREA

128 CODE AREA

**Advii**

Ibuprofen Tablets, 200 mg  
Pain Reliever/Fever Reducer (NSAID)



Caplets

24 Coated Caplets\*  
\*Capsule-Shaped Tablets

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

000067201

**Drug Facts (continued)**  
**Other information** read all warnings and directions before use. Keep carton stored at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F).

**Inactive ingredients** acetylated monoglycerides, colloidal silicon dioxide, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical grade, pharmaceutical ink, polydioxane, preparticized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

**Questions or comments?** call toll free 1-800-88-ADVIL