

Drug Facts

Active ingredient (in each tablet)

Sennosides 25 mg..... Stimulant laxative

Purpose

Uses • relieves occasional constipation (irregularity) • generally produces bowel movement in 6 to 12 hours

Warnings

Do not use laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • swallow tablet(s) with a glass of water • swallow tablet(s) whole, do not crush, break or chew

adults and children 12 years of age and older	2 tablets once or twice daily
children 6 to under 12 years of age	1 tablet once or twice daily
children under 6 years of age	ask a doctor

Drug Facts (continued)

Other information • each tablet contains: calcium 40 mg, magnesium 5 mg

• store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients acacia, calcium carbonate, carnauba wax, corn starch, dibasic calcium phosphate, FD&C blue #1 aluminum lake, iron oxide black, magnesium stearate, methylparaben, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, propylene glycol, propylparaben, shellac, silicon dioxide, sodium benzoate, sodium lauryl sulfate, sucrose, talc, titanium dioxide

Questions or comments? call 1-855-221-5432