

00201995



EXCEDRIN

PM HEADACHE

Acetaminophen 250 mg, Aspirin 250 mg (NSAID)
and Diphenhydramine Citrate 38 mg
Pain Reliever / Nighttime Sleep-Aid

100
CAPLETS

Caffeine-Free
Non-Habit Forming
ACTUAL SIZE

HALEON



00201995



Distributed by: Haleon, Warren, NJ 07059
©2023 Haleon or license; Trademarks are owned by or licensed
to the Haleon group of companies. 62000000201995

HALEON

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER SAFETY SEAL UNDER CAP IMPRINTED WITH
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts**Active ingredients (in each caplet)** **Purposes**

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Diphenhydramine citrate 38 mg.....Nighttime sleep-aid
*nonsteroidal anti-inflammatory drug

Uses • for the temporary relief of occasional headaches and
minor aches and pains with accompanying sleeplessness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering
from chicken pox or flu-like symptoms should not use this product.
When using this product, if changes in behavior with nausea and
vomiting occur, consult a doctor because these symptoms could be an
early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions.
Symptoms may include: • skin reddening • blisters • rash
if a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may
include: • hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver
damage may occur if you take

• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which
may cause severe stomach bleeding. The chance is higher if you
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs
(aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Do not use • if you have ever had an allergic reaction to
acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or
nonprescription), if you are not sure whether a drug contains
acetaminophen, ask a doctor or pharmacist.
• with any other product containing diphenhydramine, even one used
on skin
• in children under 12 years of age

Drug Facts (continued)

Ask a doctor before use if • you have liver disease
• stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney
disease
• you are taking a diuretic • you have asthma • you have glaucoma
• you have a breathing problem such as emphysema or chronic bronchitis
• you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking
a prescription drug for: • diabetes • gout • arthritis
• any other drug, or are under a doctor's care for any serious condition
• any product that contains aspirin, acetaminophen, or any other pain
reliever/fever reducer
• sedatives or tranquilizers

When using this product
• drowsiness will occur • avoid alcoholic drinks
• do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better
• sleeplessness persists continuously for more than 2 weeks. Insomnia
may be a symptom of a serious underlying medical illness.
• pain gets worse or lasts for more than 10 days
• painful area is red or swollen
• ringing in the ears or a loss of hearing occurs
• any new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It
is especially important not to use aspirin at 20 weeks or later in
pregnancy unless definitely directed to do so by a doctor because it may
cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control
Center right away. Quick medical attention is critical for adults as well
as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed
• do not use in children under 12 years of age
• adults and children 12 years of age and over: take 2 caplets at
bedtime, with a full glass of water

Drug Facts (continued)

• do not take more than 2 caplets in 24 hours, unless directed
by a doctor

Other information

• each caplet contains 75 mg calcium
• store at 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for
important information.

Inactive ingredients

benzoic acid, calcium carbonate, FD&C Blue no. 1 aluminum
lake, ferrous oxide yellow, hydroxypropyl cellulose, iron substituted
hydroxypropyl cellulose, magnesium stearate, malto-dextrin,
medium chain triglycerides, polydextrose, poly sorbate 80,
povidone, pregelatinized corn starch, silicon dioxide, titanium
dioxide, stearic acid, talc, titanium dioxide, zinc stearate

Questions? 1-800-488-7746

EXCEDRIN

PM HEADACHE

Acetaminophen 250 mg, Aspirin 250 mg (NSAID)
and Diphenhydramine Citrate 38 mg
Pain Reliever / Nighttime Sleep-Aid

ACTUAL SIZE

100
CAPLETS

Caffeine-Free
Non-Habit Forming