

00218658



EXCEDRIN

TENSION HEADACHE

ASPIRIN-FREE

Acetaminophen 500 mg, Caffeine 65 mg

Pain Reliever / Pain Reliever Aid

100
CAPLETS



ACTUAL SIZE



00218658

LOT:
EXP:



©2025 Haleon or licensor. Trademarks are owned by or licensed to the Haleon group of companies. 62000000218658
Distributed by: Haleon, Warren, NJ 07059

HALEON

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER SAFETY SEAL UNDER CAP IMPRINTED WITH
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet) **Purposes**
Acetaminophen 500 mg Pain reliever
Caffeine 65 mg Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches

Warnings

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Drug Facts (continued)

Stop use and ask a doctor if

- any new symptoms occur
 - painful area is red or swollen
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed**
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 6 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized corn starch, stearic acid, talc, titanium dioxide

Questions or comments? 1-855-297-3031

EXCEDRIN

TENSION HEADACHE

ASPIRIN-FREE

Acetaminophen 500 mg, Caffeine 65 mg
Pain Reliever / Pain Reliever Aid

ACTUAL SIZE



100
CAPLETS

HALEON

EXCEDRIN

TENSION HEADACHE

ASPIRIN-FREE

Acetaminophen 500 mg, Caffeine 65 mg
Pain Reliever / Pain Reliever Aid

100
CAPLETS