

Drug Facts (continued)

Inactive ingredients

benzalkonium chloride, dextrose, microcrystalline cellulose, phenylethyl alcohol, polysorbate 80, purified water, sodium carboxymethylcellulose

Questions or comments?

call toll-free 1-844-FLONASE
(1-844-356-6273)

FLONASE
ALLERGY RELIEF

**What problems can
FLONASE Allergy
Relief help with?**

**FLONASE Allergy Relief
helps relieve a broad
range of uncomfortable
symptoms like congestion
and itchy eyes.**



**Nasal
symptoms**



**Eye
symptoms**

**These symptoms can be
triggered by allergens
like pollen, mold, dust
or pet dander.**



**Outdoor
allergens**



**Animal
allergens**



**Indoor
allergens**

Dist. by: Haleon, Warren, NJ 07059
Made in Canada
©2024 Haleon or licensor.
Trademarks are owned by or licensed to
the Haleon group of companies.

**IMPORTANT – Peel here for complete
Drug Facts label. Children 4-11: do not
use for more than 2 months a year.**

FLONASE
ALLERGY RELIEF

Be sure to read the Quick Start Guide and
Question & Answer Leaflet inside package

Drug Facts

Active ingredient (in each spray)

Fluticasone propionate
(glucocorticoid)* 50 mcg.....reliever
*read the Question & Answer Leaflet

Purpose

TAMPER-EVIDENT features for your protection.
The product is packaged in a sealed container.
Under the cap and nozzle, each bottle has an
aluminum seal around bottle
neck. Do not use if any of these
features are torn or damaged.



209072

Drug Facts (continued)

Uses

Temporarily relieves these symptoms of hay
fever or other upper respiratory allergies:

- nasal congestion
- itchy nose
- runny nose
- itchy, watery eyes
- sneezing

Warnings

Only for use in the nose. Do not spray into
your eyes or mouth.

Do not use

- in children under 4 years of age
- to treat asthma
- if you have an injury or surgery to your
nose that is not fully healed
- if you have ever had an allergic reaction to
this product or any of the ingredients

**Ask a doctor before use if you have or
had glaucoma or cataracts**

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking

- medicine for HIV infection (such as
ritonavir)
- a steroid medicine for asthma, allergies
or skin rash
- ketoconazole pills (medicine for fungal
infection)

When using this product

- the growth rate of some children may
be slower
- stinging or sneezing may occur for a few
seconds right after use
- do not share this bottle with anyone else
as this may spread germs
- remember to tell your doctor about all the
medicines you take, including this one

Stop use and ask a doctor if

- you have, or come into contact with
someone who has, chicken pox, measles
or tuberculosis

Drug Facts (continued)

- your symptoms do not get better
within 7 days of starting use or you get
new symptoms such as severe facial
pain or thick nasal discharge. You may
have something more than allergies,
such as an infection.
- you get a constant whistling sound
from your nose. This may be a sign of
damage inside your nose.
- you get an allergic reaction to this
product. Seek medical help right away.
- you get new changes to your vision
that develop after starting this product
- you have severe or frequent
nosebleeds

If pregnant or breast-feeding, ask
a health professional before use.

Keep out of reach of children. In case
of overdose, get medical help or contact
a Poison Control Center right away.

Drug Facts (continued)

Directions

- read the Quick Start Guide for how to:
 - prime the bottle
 - use the spray
 - clean the spray nozzle
- shake gently before each use
- use this product only once a day
- do not use more than directed

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- Week 1 – use 2 sprays in each nostril
once daily
- Week 2 through 6 months – use 1 or
2 sprays in each nostril once daily, as
needed to treat your symptoms
- After 6 months of daily use – ask your
doctor if you can keep using

CHILDREN 4 TO 11 YEARS OF AGE

- the growth rate of some children may
be slower while using this product.

Drug Facts (continued)

**Children should use for the shortest
amount of time necessary to achieve
symptom relief. Talk to your child's
doctor if your child needs to use the
spray for longer than two months
a year.**

- an adult should supervise use
- use 1 spray in each nostril once daily

CHILDREN UNDER 4 YEARS OF AGE

• do not use

Other information

- you may start to feel relief the first day
and full effect after several days of
regular, once-a-day use
- store between 4° and 30°C (39°
and 86°F)
- keep this label and enclosed materials.
They contain important additional
information.