

Drug Facts (continued)

- keep this label and enclosed materials. They contain important additional information.

Inactive ingredients

benzalkonium chloride solution, carboxymethylcellulose sodium, dextrose anhydrous, edetate disodium, microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

call toll-free 1-844-FLONASE
(1-844-356-6273)

FLONASE
SENSIMIST ALLERGY RELIEF

What problems can
**FLONASE SENSIMIST
Allergy Relief help with?**

**FLONASE SENSIMIST
Allergy Relief helps
relieve a broad range of
uncomfortable symptoms
like congestion and
itchy eyes*.**



**Nasal
symptoms**



**Eye
symptoms***

*for ages 12 and older.

These symptoms can be
triggered by allergens
like pollen, mold, dust
or pet dander.



**Outdoor
allergens**



**Animal
allergens**



**Indoor
allergens**

Dist. by: Haleon, Warren, NJ 07059
Made in Singapore
©2024 Haleon or licensors.
Trademarks are owned by or licensed to
the Haleon group of companies.

IMPORTANT – Peel here for complete
Drug Facts label. Children 2-11: do not
use for more than 2 months a year.

FLONASE
SENSIMIST ALLERGY RELIEF

Be sure to read the Quick Start Guide and
Question & Answer Leaflet inside package

Drug Facts

Active ingredient (in each spray)	Purpose
Fluticasone furoate (glucocorticoid)* 27.5 mcg.....reliever	Allergy symptom
*read the Question & Answer Leaflet	

TAMPER-EVIDENT features for your protection. The
product is packaged in a sealed container. The bottle
is housed within a plastic dispenser that prevents direct
access to the bottle. Do not use if the
container or any part of the plastic
dispenser, including the dispenser
bottom, is damaged or broken.



209081

Data Matrix Code #209081

Drug Facts (continued)

Uses

Temporarily relieves these symptoms of hay
fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose
- itchy, watery eyes (for ages 12 and older)

Warnings

Only for use in the nose. Do not spray into
your eyes or mouth.

Do not use

- in children under 2 years of age
- to treat asthma
- if you have an injury or surgery to your
nose that is not fully healed
- if you have ever had an allergic reaction to
this product or any of the ingredients

Ask a doctor before use if you have or
had glaucoma or cataracts

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking

- medicine for HIV infection (such as
ritonavir)
- a steroid medicine for asthma, allergies
or skin rash
- ketoconazole pills (medicine for fungal
infection)

When using this product

- the growth rate of some children may
be slower
- stinging or sneezing may occur for a few
seconds right after use
- do not share this bottle with anyone else
as this may spread germs
- remember to tell your doctor about all the
medicines you take, including this one

Stop use and ask a doctor if

- you have, or come into contact with
someone who has, chicken pox, measles
or tuberculosis

Drug Facts (continued)

- your symptoms do not get better
within 7 days of starting use or you get
new symptoms such as severe facial
pain or thick nasal discharge. You may
have something more than allergies,
such as an infection.
- you get a constant whistling sound
from your nose. This may be a sign of
damage inside your nose.
- you get an allergic reaction to this
product. Seek medical help right away.
- you get new changes to your vision
that develop after starting this product
- you have severe or frequent
nosebleeds

If pregnant or breast-feeding, ask
a health professional before use.

Keep out of reach of children. In case
of overdose, get medical help or contact
a Poison Control Center right away.

Drug Facts (continued)

Directions

- read the Quick Start Guide for how to:

- shake vigorously before each use
- prime the bottle
- use the spray
- sniff gently after each spray
- clean the spray nozzle with a clean dry
tissue
- use this product only once a day
- do not use more than directed

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- Week 1 – use 2 sprays in each nostril
once daily
- Week 2 through 6 months – use 1 or
2 sprays in each nostril once daily, as
needed to treat your symptoms
- After 6 months of daily use – ask your
doctor if you can keep using

Drug Facts (continued)

CHILDREN 2 TO 11 YEARS OF AGE

- the growth rate of some children may
be slower while using this product.
**Children should use for the shortest
amount of time necessary to achieve
symptom relief. Talk to your child's
doctor if your child needs to use the
spray for longer than two months
a year.**

- an adult should supervise use
- use 1 spray in each nostril once daily

CHILDREN UNDER 2 YEARS OF AGE

- do not use

Other information

- you may start to feel relief the first day
and full effect after several days of
regular, once-a-day use
- store between 15° - 30°C (59° - 86°F)
- do not refrigerate or freeze