

Purposes

Drug Facts

Active ingredients (per tablet)

Calcium carbonate 750 mgAntacid

Simethicone 250 mg..... Uses for the relief of • pressure, bloating, and fullness commonly referred to as gas • gas associated with heartburn, sour stomach or acid indigestion acid indigestion
beartburn
sour stomach
upset stomach

associated with these symptoms Warnings

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs. When using this product • do not exceed 2 tablets in 24 hours except under the advice and supervision of a doctor • do not use the maximum dosage for more than 2 weeks, except under the advice and supervision of a doctor

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. adults and children 12 years and older: chew 1 or 2 tablets

as symptoms occur, or as directed by a doctor. Chew or crush tablets completely before swallowing. • do not take for symptoms that persist for more than 2 weeks, unless advised by a doctor

Other information • each tablet contains: elemental calcium 295 mg store in a cool, dry place at no higher than 25°C (77°F)

contains FD&C Yellow 5 Lake (tartrazine) as a color additive

Inactive inaredients alcohol, ammonium hydroxide, beeswax, carnauba wax, citric acid, corn syrup, dextrin, ethyl acetate, FD&C red 40, FD&C red 40 lake, FD&C vellow 5 lake (tartrazine), FD&C vellow 6, FD&C vellow 6 lake, flavors, gum arabic, isopropyl alcohol, maltodextrin, methylparaben, n-butyl alcohol,

pharmaceutical ink, phosphoric acid, propylene glycol, propylparaben, purified water, shellac, sodium benzoate, sorbic acid, sorbitol, soy lecithin, sucrose, titanium dioxide, vegetable oil (coconut origin)

Questions or comments? 1-855-328-5204