Drug Facts Purposes Active ingredients (in each 5 mL / teaspoonful) 0 Aluminum hydroxide 254 mg......Antacid O. Magnesium carbonate 237.5 mg......Antacid Uses relieves •heartburn acid indigestion sour stomach

00213322

Warnings

Ask a doctor or pharmacist before use if you

upset stomach associated with these symptoms





Drug Facts (continued)

have kidney disease

 are on a sodium-restricted diet or a magnesium-restricted diet are taking a prescription drug. Antacids may

interact with certain prescription drugs. When using this product

 do not take more than 16 teaspoonfuls in 24 hours

 do not use the maximum dosage for more than 2 weeks, except under the advice and supervision of a doctor · laxative effect may occur

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- shake well • take 2-4 teaspoonfuls four times a day or as
- directed by a doctor take after meals and at bedtime
- · dispense product only by spoon or other measuring device

Drug Facts (continued)

Other information each teaspoon (5 mt.) contains: magnesium 75 mg, sodium 11 mg store at up to 25°C (77°F). Avoid freezing.

keep tightly closed

Inactive ingredients benzyl alcohol, edetate disodium, glycerin, natural and artificial flavors, purified water, simethicone emulsion. sodium alginate, sodium saccharin, sorbitol

Questions or comments? 1-800-452-0051

solution, xanthan gum