



NicoDerm CQ is a three step program. You start with the highest level of medicine and gradually step down your dose. If you smoke 10 or less cigarettes per day, start with Step 2.

NDC 0135-0196-02



**Nicotine Transdermal System STOP SMOKING AID** STEP 3 14 clear patches (2-week kit)

Not for sale to those under 18 years of age. Proof of age required.

• Not for sale in vending machines or from any source where proof of age cannot be verified.

Trademarks are owned by or licensed to the GSK group of companies.

For more information and for a FREE individualized stop smoking program, please visit www.nicodermcq.com or see inside for more details.

U.S. Patent No. 7,622,136 8.075.911

## NicoDerm Nicotine Transdermal System 7 mg Delivered over 24 Hours STOP SMOKING AID



SMARTCONTROL TECHNOLOGY

STEP C

14 clear patches (2-week kit)

FOR USE AFTER

**COMPLETING STEP 2** 

7 mg



Nicotine Transdermal System **STOP SMOKING AID** STEP 3 14 clear patches (2-week kit)

What is the NicoDerm CQ Patch and How is it Used? NicoDerm CQ is a small, nicotine-containing patch. When you put on a NicoDerm CQ patch, nicotine passes through the skin and into your body. NicoDerm CQ is very thin and uses special material to control how fast nicotine passes through the skin. Unlike the sudden jolts of nicotine delivered by cigarettes, the amount of nicotine you receive remains relatively smooth throughout the 16 or 24 hour period you wear the NicoDerm CQ patch. This helps to reduce cravings you may have for nicotine.



## 200699 200699 200699 200699

ACTUAL

SIZE

STEP 3 14 clear patches

(2-week kit)

gsk



Read carton and enclosed User's Guide before using this product. Keep this carton and User's Guide. They contain important information		
NicoDerm	Drug Facts	Drug Facts (continued)
CQ TO INCREASE YOUR	Active ingredient (in each patch) Purpose Nicotine, 7 mg delivered over 24 hoursStop smoking aid	<ul> <li>begin using the patch on your quit day</li> <li>if you smoke more than 10 cigarettes per day, use according to the following 10-week schedule:</li> </ul>
<b>SUCCESS IN QUITTING:</b> 1. You must be motivated to quit.	<b>Use</b> reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking	STEP 1         STEP 2         STEP 3           Use one 21 mg patch/day         Use one 14 mg patch/day         Use one 7 mg patch/day
<ol> <li>Complete the full treatment program, applying a new patch every day.</li> </ol>	Warnings If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine	patch/day         patch/day         patch/day           Weeks 1-6         Weeks 7-8         Weeks 9-10           • If you smoke 10 or less cigarettes per day, do not use         STEP 1 (21 mg). Start with STEP 2 (14 mg) for 6 weeks, then STEP 3 (7 mg) for 2 weeks and then stop.
<ol> <li>Use with a support program as described in the enclosed User's Guide.</li> </ol>	is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.	<ul> <li>steps 2 and 3 allow you to gradually reduce your level of nicotine. Completing the full program will increase your</li> </ul>
For your family's protection, NicoDerm <i>CQ</i> patches are supplied in child resistant pouches. Do not use if individual pouch is open or torn.	Ask a doctor before use if you have • heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. • high blood pressure not controlled with medication. Nicotine can increase your blood pressure. • an allergy to adhesive tape or have skin problems because you are more likely to get rashes • stomach ulcer or diabetes • history of seizures	<ul> <li>chances of quitting successfully.</li> <li>apply one new patch every 24 hours on skin that is dry, clean and hairless. Save pouch for disposing of the patch after use.</li> <li>remove backing from patch and immediately press onto skin.</li> <li>Hold for 10 seconds.</li> <li>wash hands after applying or removing patch. Throw away the patch by folding sticky ends together. Replace in its pouch and discard. See enclosed User's Guide for safety and handling.</li> </ul>
	Ask a doctor or pharmacist before use if you are • using a non-nicotine stop smoking drug • taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.	<ul> <li>you may wear the patch for 16 or 24 hours</li> <li>if you crave cigarettes when you wake up, wear the patch for 24 hours</li> <li>if you have vivid dreams or other sleep disturbances, you may remove the patch at bedtime and apply a new one in</li> </ul>
Distributed by: GSK Consumer Healthcare	When using this product • if you have vivid dreams or other sleep disturbances remove this patch at bedtime	the morning • the used patch should be removed and a new one applied to a different skin site at the same time each day • do not wear more than one patch at a time • do not used the time of the same time second the same • do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours • it is important to complete treatment. If you feel you need to
Warren, NJ 07059 Made in Ireland ©2022 GSK group of companies or	Stop use and ask a doctor if • skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash • irregular heartbeat or palpitations occur • you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, weakness and rapid heartbeat • you have symptoms of an allergic reaction (such as difficulty breathing or rash)	
its licensor.		use the patch for a longer period to keep from smoking, talk to your health care provider.
	Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Dispose of the used patches by folding sticky ends together. Replace in pouch and discard.	<b>Other information</b> • store at 20-25°C (68-77°F)
		Inactive ingredients ethylene vinyl acetate-copolymer, polyisobutylene and high density polyethylene between clear polyester backings
	Directions • if you are under 18 years of age, ask a doctor before use • before using this product, read the enclosed User's Guide for complete directions and other information	Questions or comments? call toll-free 1-800-834-5895
	Includes User's Guide with My	Quit Program Enrollment Offer.

200699

