




Opening Directions:
Squeeze child resistant tabs on each side.
Flip up the top of vial. Turn upside down
and shake to remove lozenge.



Drug Facts (continued)

- nicotine lozenge is a medicine and must be used a certain way to get the best results
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- store between 20°-25°C (68°-77°F)
- keep the vial tightly closed and protect from light

Inactive ingredients
acesulfame potassium, anhydrous dibasic calcium phosphate, crospovidone, FD&C blue no. 2 aluminum lake, flavors, hypromellose, magnesium stearate, mannitol, microcrystalline cellulose, sodium bicarbonate, sodium carbonate anhydrous, sucralose, xanthan gum, xylitol

Questions or comments?
Call 1-855-297-3031

Distributed by:
Haleon
Warren, NJ 07059
Made in Switzerland
Trademarks are owned by or licensed to
the Haleon group of companies.
©2025 Haleon group of companies or its licensor.
Patent Pending
221954

NDC 0135-8060-81 HALEON

NICORETTE

nicotine polacrilex lozenge, 4 mg
stop smoking aid
mini
Lozenge

4mg

FOR THOSE WHO
SMOKE THEIR
FIRST CIGARETTE
WITHIN 30 MINUTES
OF WAKING UP.

If you smoke your
first cigarette
MORE THAN 30 MINUTES
after waking up, use
Nicorette 2 mg Lozenge.



81 LOZENGES,
4 mg EACH (3 Vials of 27)

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. **Use Enough** — Use at least 9 Nicorette mini Lozenges per day during the first six weeks.
3. **Use Long Enough** — Use Nicorette mini Lozenges for the full 12 weeks.
4. **Use with a support program** as directed in the enclosed User's Guide.

■ not for sale to those under 18 years of age

■ proof of age required

■ not for sale in vending machines or from any source where proof of age cannot be verified

TAMPER EVIDENT FEATURE:
Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken. Retain outer carton for full product uses, directions and warnings.

For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.



Drug Facts

Active ingredient (in each lozenge)	Purpose
Nicotine polacrilex, 4 mg.....	Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours