parodontax ACTIVE GUM HEALTH

Drug Facts		
Active ingredient Purpose Cetylpyridinium chloride 0.07%w/wAntigingivitis, Antiplaque		
Uses helps prevent and reduce plaque that leads to ■ gingivitis, an early form of gum disease ■ bleeding gums		
Warnings Stop use and ask a dentist if irritation occurs gingivitis, bleeding, or redness persists for more than 2 weeks you have painful or swollen gums, pus from the gum line, loose teeth, or increasing spacing between the teeth. These may be signs or symptoms of periodontitis, a serious form of gum disease. Keep out of reach of children under 6 years of age. If more than used for rinsing is		
Diffection is a datust and children by ears of age and older: measure twice up to 10 millitters mark inside of cap. Vigorously swish 20 millitters of the rinse between your teeth twice a day for 30 seconds and then spit out. Do not swallow the rinse. Do not rinse with water. Do not drink from the bottle. ■ children 6 years of age: do not use		3.75 "
Other information ■ this rinse is not intended to replace brushing or flossing ■ may cause staining ■ Do not store above 25°C or 77°F		
Inactive ingredients Water, glycerin, flavor, poloxamer 188, sodium saccharin, propylene glycol, sodium benzoate, sucralose, benzoic acid		
Questions or comments? 1-855-328-5202		
*Kills plaque bacteria associated with gingivitis and odor-causing bacteria in a laboratory test Do not use if the printed seal on the cap is broken or missing. ALWAYS FOLLOW THE LABEL Distributed by: GSK Consumer Healthcare Warren, NJ 07059 Trademarks owned by or licensed by GSK. ©2021 GSK or licensor. 1.855-328-5202 1.855-328-5202 1.855-328-5202		
	Active ingredient Purpose Cetylpyridinium chloride 0.07%w/w. Antigingivitis, Antiplaque USeS helps prevent and reduce plaque that leads to gingivitis, an early form of gum disease bleeding gums gingivitis, an early form of gum disease bleeding gums Stop use and ask a dentist if iritation occurs gingivitis, bleeding, or redness persists for more than 2 weeks you have painful or swollen gums, pus from the gum line, loose teeth, or increasing spacing between the teeth. These may be signs or symptoms of periodonitiis, a serious form of gum disease. Keep out of reach of children under 6 years of age and older: measure twice up to 10 milliliters mark inside of cap. Vigorously swish 20 milliliters of the rinse between he teeth with earls on ot drink from the bottle. Directions and your obseconds and then spit out. Do not swallow the rinse. Do not milliliters of age: and older: measure twice up to 10 milliliters are inside of cap. Vigorously swish 20 milliliters of the rinse between your teeth twice a day for 30 seconds and then spit out. Do not swallow the rinse. Do not not rinse with water. Do not drink from the bottle. Other information this rinse is not intended to replace brushing or flossing may cause staining Do not store above 25°C or 77°F Inactive ingredients Vater, glycerin, flavor, poloxamer 188, sodium saccharin, propylene glycol, sodium benzoate, sucralose, benzoic acid Questions or comments? 1-855-328-5202 rKlils plaque bacteria associated with gingivitis and odor-causing bacteria in a laboratory test Waren, NU 0705 LO020007BB Is broken o	Active ingredient Purpose Cetylpyridinium chloride 0.07%w/w. Antigingivitis, Antiplaque USeS helps prevent and reduce plaque that leads to gingivitis, an early form of gum disease of gum disease bleeding gums gingivitis, bleeding, or redness persists for more than 2 weeks you have painful or swollen gums, pus from the gum line, loose techt, or increasing spacing between the techt. These may be signs or symptoms of periodontitis, a serious form of gum disease. Keep out of reach of children under 6 years of age and older: measure twice up to 10 milliliters mark inside of cap. Ugorusly swish 20 milliliters of the rinse between the techt or increasing and older: measure twice up to 10 milliliters of the signs or supervise use Directions = adults and children Gyears of age and older: measure twice up to 10 milliliters of the rinse between your teeth twice a day for 30 seconds and then spit out. Do not swallow the rinse. Do not not milliliters of the rinse between your teeth wider. Do not drink from the bottle. — children 6 years of age: do not use Other information = this rinse is not intended to replace brushing or flossing = may cause staining = Don to store above 25°C or 77°F Inactive ingredients Nater, glycerin, flavor, poloxamer 188, sodium saccharin, propylene glycol, sodium benzoate, sucralose, benzoic acid Questions or comments? 1-855-328-5202 fklis plaque bacteria associated with gingivitis and odor-causing bacteria in a laboratory test bor tore by cits broken or missing. LO