

Drug Facts

Active ingredient

Cetylpyridinium chloride 0.07% w/w.....Antigingivitis, Antiplaque

Purposes

Uses helps prevent and reduce plaque that leads to ■ gingivitis, an early form of gum disease ■ bleeding gums

Warnings

Stop use and ask a dentist if ■ irritation occurs ■ gingivitis, bleeding, or redness persists for more than 2 weeks ■ you have painful or swollen gums, pus from the gum line, loose teeth, or increasing spacing between the teeth. These may be signs or symptoms of periodontitis, a serious form of gum disease.

Keep out of reach of children under 6 years of age. If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ **adults and children 6 years of age and older:** vigorously swish 20 milliliters (4 teaspoonfuls) of rinse between your teeth twice a day for 30 seconds and then spit out. Do not swallow the rinse. Do not rinse with water. Do not drink from the bottle. ■ **children 6 years to under 12 years of age:** supervise use ■ **children under 6 years of age:** do not use

Other information ■ this rinse is not intended to replace brushing or flossing ■ may cause staining ■ do not store above 25°C (77°F) ■ this product may turn cloudy. Its antimicrobial properties are not affected.

Inactive ingredients water, glycerin, propylene glycol, poloxamer 407, flavor, benzoic acid, sodium benzoate, sodium saccharin, sucralose, sodium hyaluronate, blue 1

Questions or comments? 1-866-844-2797

ALWAYS FOLLOW THE LABEL