

# Drug Facts

## Active ingredient

Sodium fluoride 0.05% (0.02% w/v fluoride ion)..... Anticavity

## Purpose

## Uses

- aids in the prevention of dental cavities.
- the combined daily use of a fluoride preventive treatment rinse and a fluoride toothpaste can help reduce the incidence of dental cavities.

## Warning

**Keep out of reach of children.** If more than used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away.

## Directions

- **adults and children 6 years of age and older:** use once a day after brushing your teeth with a toothpaste.
- pour 10 milliliters into cap (10 mL mark on inside of cap); do not fill above the 10 mL mark.
- vigorously swish 10 milliliters of rinse between your teeth for 1 minute and then spit out.
- do not swallow the rinse.
- do not drink from the bottle.
- do not rinse mouth with water.
- do not eat or drink for 30 minutes after rinsing.
- instruct children under 12 years of age in good rinsing habits (to minimize swallowing).
- supervise children as necessary until capable of using without supervision.
- **children under 6 years of age:** Consult a dentist or doctor.

## Other information

- do not store above 25°C (77°F)

## Inactive ingredients

water, sorbitol, sodium benzoate, poloxamer 407, benzyl alcohol, polysorbate 20, flavor, propylene glycol, sodium saccharin, disodium EDTA, disodium phosphate, monosodium phosphate, calcium disodium EDTA, cetylpyridinium chloride, potassium sorbate, FD&C blue no. 1

## Questions or comments? 1-866-844-2797