#### Robitussin Maximum Strength Cough+Chest Congestion DM

Do not take both products within 6 hours of each other.

# Drug Facts

# Active inaredients (in each 20 mL)

Purposes

Dextromethorphan HBr. USP 20 mg......Cough suppressant Guaifenesin, USP 400 mg. .. Expectorant

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phleam (mucus) and thin bronchial secretions to drain bronchial tubes

# Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression. psychiatric, or emotional conditions, or Parkinson's disease). or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- cough that occurs with too much phleam (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

## Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product mmL = milliliter
- this adult product is not intended for use in children under 12 years of ane

TE yours or ago	
age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 19 vegre	do not usa

# Other information

■ each 20 mL contains: sodium 13 mg store at 20-25°C (68-77°F). Do not refrigerate.

**Robitussin Maximum Strength** Cough+Chest Congestion DM (continued)

#### Drug Facts (continued)

Inactive inaredients anhydrous citric acid. carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40. glycerin, liquid glucose, menthol, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose, triacetin, xanthan qum

**Questions or comments?** call weekdays from 8 AM to 6 PM FST at 1-800-245-1040

## **Robitussin Maximum Strength** Nighttime Cough DM

Do not take both products within 6 hours of each other.

# Drug Facts

Active ingredients (in each 20 mL) Purposes

Dextromethorphan HBr, USP 30 mg....... Cough suppressant Doxylamine succinate LISP 12.5 mg

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
  sneezing
  itchy, watery eyes
- itching of the nose or throat
- controls the impulse to cough to help you sleep

# Warnings

## Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric. or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

# Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- a cough that occurs with too much phleam (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

**Robitussin Maximum Strength** Nighttime Cough DM (continued)

# Drug Facts (continued)

When using this product do not use more than directed

- marked drowsiness may occur avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

# **Directions** ■ measure only with dosing cup provided

- keep dosing cup with product mL = milliliter do not take more than 4 doses in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 6 hours
children under 12 years	do not use

#### Other information

each 20 mL contains: sodium 14 mg ■ store at 20-25°C (68-77°F)

Inactive inaredients anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, glycerin, liquid glucose, menthol, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water sodium benzoate sodium citrate sucralose triacetin xanthan gum

**Questions or comments?** call weekdays from 9 AM to 5 PM FST at 1-800-245-1040

# TAMPER-EVIDENT INNER UNITS:

Do Not Use if Tamper-Evident feature on either bottle cap is broken or missing. See individual inner cartons and bottle labels for specific Tamper-Evident features

©2024 Haleon or licensor. Pat. Info www.productpats.com See individual boxes for foreign country of origin.

FAST, EFFECTIVE cough relief

MAX

DAY&NIGHT

Robitussin

Cough+Chest Nighttime Congestion DM Cough DM

**ADULT** 

**MAXIMUM STRENGTH** 

DEXTROMETHORPHAN HBr (Cough Suppressant)

✓ Relieves Chest Congestion

√ Thins & Loosens Mucus

GUAIFENESIN (Expectorant)

✓ Controls Cough

DEXTROMETHORPHAN HBr (Cough Suppressant) DOXYLAMINE SUCCINATE (Antihistamine)

**HALEON** 

- √ Controls Cough
- √ Relieves Runny Nose & Sneezina

**MAXIMUM STRENGTH** 



For Ages 12 & Over 4 FL OZ (118 mL)

# Robitussin





www.StopMedicineAbuse.org