

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking any other pain reliever/fever reducer
- taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.

Directions

- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage.
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

Drug Facts (continued)

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 12 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, artificial flavor, edetate disodium, FD&C red no. 40, glycerin, menthol, polyethylene glycol, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

STRONG SOOTHING ACTION
and powerful relief for your worst symptoms

Also available:



Distributed by: Haleon, Warren, NJ 07059
For most recent product information, visit www.robitussin.com
Trademarks owned or licensed by Haleon.
Pat. Info www.productpats.com
©2023 Haleon or licensor. Made in Canada



ADULT NEW FORMULA HALEON

Robitussin

MAXIMUM STRENGTH

SEVERE MULTI-SYMPTOM
Cough Cold + Flu

Nighttime

ACETAMINOPHEN (Pain Reliever/Fever Reducer)
DIPHENHYDRAMINE HCl (Antihistamine/Cough Suppressant)

Relieves Cough to Help You Rest

- ✓ Cough, Sore Throat
- ✓ Head & Body Aches, Fever
- ✓ Runny Nose, Sneezing

FAST, POWERFUL
multi-symptom relief

CF
NIGHTTIME
MAX

For Ages 12 & Over 4 FL OZ (118 mL)

Do Not Use if breakable ring on bottle cap is separated or missing.

Drug Facts

Active ingredients
(in each 20 mL)

Purposes

Acetaminophen 650 mgPain reliever/Fever reducer
Diphenhydramine HCl 25 mgAntihistamine/
Cough suppressant

Uses

- temporarily relieves these symptoms occurring with a cold or flu, hay fever, or other respiratory allergies:
 - cough due to minor throat and bronchial irritation
 - headache
 - sore throat
 - minor aches and pains
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily relieves your cough to help you rest
- temporarily reduces fever

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 6 doses in any 24-hour period, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to sedate a child or to make a child sleepy
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin