

Children's Robitussin Cough & Chest Congestion DM

Do not take both products within 6 hours of each other.

Drug Facts

Active ingredients (in each 20 mL) Purposes
 Dextromethorphan HBr 20 mg Cough suppressant
 Guaifenesin 200 mg Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter

age	dose
children under 4 years	do not use
children 4 to under 6 years	5 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
adults and children 12 years and over	20 mL every 4 hours

Other information

- each 20 mL contains: sodium 21 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Children's Robitussin Cough & Chest Congestion DM

(continued)

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, glycerin, liquid glucose, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

Children's Robitussin Nighttime Cough Long-Acting DM

Do not take both products within 6 hours of each other.

Drug Facts

Active ingredients (in each 10 mL) Purposes
 Chlorpheniramine maleate, USP 2 mg Antihistamine
 Dextromethorphan HBr, USP 15 mg Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

- When using this product**
- do not use more than directed
 - marked drowsiness may occur
 - avoid alcoholic drinks
 - alcohol, sedatives, and tranquilizers may increase drowsiness

Children's Robitussin Nighttime Cough Long-Acting DM

(continued)

Drug Facts (continued)

- be careful when driving a motor vehicle or operating machinery
 - excitability may occur, especially in children
- Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period

age	dose
children under 6 years	do not use
children 6 to under 12 years	10 mL every 6 hours
adults and children 12 years and over	20 mL every 6 hours

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, lactic acid, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? call weekdays from 8 AM to 6 PM EST at 1-800-245-1040

Distributed by: Haleon, Warren, NJ 07059
 For most recent product information, visit www.robitussin.com
 Trademarks owned or licensed by Haleon.
 ©2023 Haleon or licensor.
 Pat. Info www.productpats.com
 Some product Made in Canada. See individual boxes for foreign country of origin.

TAMPER-EVIDENT INNER UNITS:

Do Not Use if Tamper-Evident feature on either bottle cap is broken or missing. See individual inner cartons and bottle labels for specific Tamper-Evident features.

DAY & NIGHT

COUGH RELIEF

Children's

Robitussin

HALEON

AGES 4 & OVER

Cough & Chest Congestion

DM

DEXTROMETHORPHAN HBr (Cough Suppressant)
GUAIFENESIN (Expectorant)

Relieves:

- ✓ Chest Congestion/Mucus
- ✓ Cough

grape flavor

4 FL OZ (118 mL)

AGES 6 & OVER

Nighttime Cough

DM

Long-Acting

CHLORPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)

Relieves:

- ✓ Cough Up to 8 Hours
- ✓ Runny Nose

Alcohol-Free

fruit punch flavor

4 FL OZ (118 mL)

Children's Robitussin liquid is specially formulated to provide soothing action, control your child's cough -plus relieves other cold symptoms.



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org