

Do Not Use if breakable ring is separated or missing.
Packaged with Tamper-Evident bottle cap.

Robitussin

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 15 mg...Cough suppressant
Doxylamine succinate, USP 6.25 mg.....Antihistamine

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period

age	dose
children under 6 years	do not use
children 6 to under 12 years	10 mL every 6 hours
adults and children 12 years and older	20 mL every 6 hours

Other information

- each 10 mL contains: sodium 10 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, blueberry juice concentrate, carboxymethylcellulose sodium, glycerin, lactic acid, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments? call weekdays from 9 AM to 5 PM EST at 1-800-245-1040

Dist. by: Haleon, Warren, NJ 07059

©2024 Haleon or licensors.

Trademarks owned or licensed by Haleon.

For most recent product information, visit www.robitussin.com

Pat. Info www.productpats.com Made in Canada

62000000211561

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Children's

Robitussin



Nighttime Cough



Children's

Children's

Robitussin Robitussin

Honey

Honey



For Ages 6+

Nighttime Cough



DEXTROMETHORPHAN HBr
(Cough Suppressant)
DOXYLAMINE SUCCINATE (Antihistamine)

ALCOHOL
FREE

LONG-ACTING

- Relieves:
- ✓ Cough up to 8 hours
 - ✓ Runny nose

Taste the
Real Honey



4 FL OZ (118 mL)

Made with
Real Honey



- Gluten Free
- No Artificial Colors



85% FPO

3 0031 8762 12 7