

COPY AREA

NO ON INK COATING

NEW! ADULT

Robitussin

Elderberry MEDI-SOOTHERS

NEW! ADULT

HALEON

Robitussin

Elderberry



MEDI-SOOTHERS

DEXTROMETHORPHAN HBr (Cough Suppressant) 5 mg
MENTHOL (Oral Anesthetic) 5 mg

Cough + Sore Throat

Taste the Real Elderberry



Elderberry Flavor

- ✓ Relieves Cough for up to 4 HOURS
- ✓ Relieves Sore Throat Pain

For Ages 12+

16 MEDICATED LOZENGES

ADULT

Robitussin

Elderberry MEDI-SOOTHERS

Taste the Real Elderberry



Keep carton for complete information.
Do not use if blister is broken or damaged!

Distributed by: Haleon, Warren, NJ 07059
Trademarks owned or licensed by Haleon.
©2023 Haleon or its licensor.
For most recent product information, visit www.robitussin.com

202559

ADULT

Robitussin

Elderberry MEDI-SOOTHERS

Taste the Real Elderberry



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts

Active ingredients (in each lozenge)

Dextromethorphan HBr 5 mg Cough suppressant
Menthol 5 mg Oral anesthetic

Purposes

- temporarily relieves
- cough due to minor throat and bronchial irritation as may occur with the common cold
- occasional minor irritation, pain, sore mouth, and sore throat

Warnings

- Sore throat warning**
- If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

- Stop use and ask a doctor or dentist if**
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.
- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years of age and older: take 2 lozenges (one immediately after the other) every 4 hours as needed. Allow each lozenge to dissolve slowly in the mouth.
- do not take more than 12 lozenges in any 24-hour period, or as directed by a doctor
- children under 12 years of age: do not use

Other information

Store at 20°-25° C (68°-77°F)

Inactive ingredients

corn syrup, elderberry juice concentrate, glycerin, natural flavors, propylene glycol, purified water, soy lecithin, sucralose, sucrose

Questions or comments?

call weekdays from 8 am to 6 pm EST at 1-800-245-1040