Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* is broken or missing.

***Among those with a preference vs. original formulation.**

Distributed by: Haleon, Warren, NJ 07059
Trademarks are owned by or licensed to the Haleon group of companies.
©2025 Haleon group of companies or its licensor.
For most recent product information, visit www.robitussin.com

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

For Position Only
3 00310 34074 3

NEW! IMPROVED FLAVOR* HALEON

Robitussin

Nighttime Cough DM Soft CHEWS

DEXTROMETHORPHAN HBr 15 mg (Cough Suppressant)
DOXYLAMINE SUCCINATE 6.25 mg (Antihistamine)

FAST* relief *ANYWHERE
Cough, runny nose & sneezing



UP TO 8HR COUGH RELIEF

20 CHEWABLE TABLETS
Chew tablets completely before swallowing

***Berry* flavor**

Drug Facts

Active ingredients (in each chewable tablet) — Purposes

Dextromethorphan equivalent to 15 mg dextromethorphan HBr ..... Cough suppressant
Doxylamine succinate 6.25 mg..... Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if

50207846

217072

Peel Here

Drug Facts (continued)

your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than 4 doses (8 chewable tablets) in 24-hours, unless directed by a doctor.
- chew tablets completely before swallowing
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	take 2 chewable tablets every 6 hours, as needed
children under 12 years	do not use

Other information

- **each chewable tablet contains:** calcium 10 mg, potassium 10 mg, sodium 7 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

calcium gluconate monohydrate, carboxymethylcellulose calcium, carnauba wax, corn syrup solids, crospovidone, D&C red no. 30 aluminum lake, FD&C blue no. 1 aluminum lake, glycerin, mannitol, natural and artificial flavors, pregelatinized starch, simethicone, sodium gluconate, sorbitol, sorbitol solution, sucralose, sucrose, whole dry milk

Questions or comments?

call weekdays from 8 AM to 6 PM EST at **1-855-297-3031**