Drug Facts Drug Facts (continued) Active ingredient Directions Purposes · adults and children 12 years of age and older: apply at least a 1-inch strip of the product onto a soft bristle toothbrush. Uses brush teeth thoroughly for at least 1 minute twice a day (morning and evening), and not more than 3 times a aids in the prevention of dental cavities.
helps interfere with harmful effects of plaque associated with day, or as recommended by a dentist or doctor. Make sure to brush all sensitive areas of the teeth. Minimize builds increasing protection against painful sensitivity of the teeth to cold, heat, acids, sweets. ainaivitis. swallowing. Spit out after brushing. or contact. children under 12 years of age: Consult a dentist or doctor. Warnings Other information Stop use and ask a dentist if products containing stannous fluoride may produce surface staining of the teeth. Adequate toothbrushing may the problem persists or worsens. Sensitive teeth may indicate a serious problem that may need prompt care by prevent these stains which are not harmful or permanent and may be removed by your dentist. a dentist. • pain/sensitivity still persists after 4 weeks of use. • qinqivitis, bleeding, or redness persists for • this product is specially formulated to help prevent staining. • store below 30°C (86°F) more than 2 weeks. • you have painful or swollen gums, pus from the gum line, loose teeth, or increasing Inactive ingredients glycerin, PEG-8, hydrated silica, pentasodium triphosphate, flavor, sodium spacing between the teeth. These may be signs or symptoms of periodontitis, a serious form of gum disease. lauryl sulfate, titanium dioxide, polyacrylic acid, sodium saccharin, cocamidopropyl betaine Keep out of reach of children. If more than used for brushing is accidentally swallowed, get medical help or contact a Poison Control Center right away. Questions or comments? 1-866-844-2797 ALWAYS FOLLOW THE LABEL