

READ ALL WARNINGS AND DIRECTIONS ON
CARTON BEFORE USE. KEEP CARTON FOR REFERENCE.
DO NOT DISCARD.

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 1000 mg.....	Pain reliever/Fever reducer
Chlorpheniramine maleate 4 mg.....	Antihistamine
Dextromethorphan HBr 30 mg.....	Cough suppressant
Pseudoephedrine HCl 60 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - minor sore throat pain
 - runny nose
 - cough due to minor throat and bronchial irritation
 - nasal and sinus congestion
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- Do not use** • in a child under 12 years of age
- if you are allergic to acetaminophen
 - with any other drug containing acetaminophen (prescription or nonprescription).
If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

Drug Facts (continued)

- cough comes back or occurs with rash or headache that lasts
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- measure the dose correctly using the enclosed dosing cup
- adults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persist
- do not take more than 3 doses (90 mL) in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	30 mL every 6 hours
children under 12 years of age	do not use

Other information

- each 30 mL contains: sodium 26 mg
- store at 20-25°C (68-77°F). Do not refrigerate. Keep bottle tightly closed.

Inactive ingredients alcohol, anhydrous citric acid, carboxymethylcellulose sodium, D&C red no. 33, FD&C red no. 40, glycerin, natural and artificial flavors, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

Questions or comments? 1-855-328-5259

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

**DO NOT USE IF TEAR BAND ON CAP IS BROKEN
OR MISSING.**

Distributed by: Haleon
Warren, NJ 07059

Made in Canada

©2024 Haleon group of companies or its licensor.
Trademarks are owned by or licensed to the
Haleon group of companies. 62000000209276
Pat. Info www.productpats.com

**FLU RELIEF
MAX STRENGTH***

**NASAL
DECONGESTANT**

NIGHTTIME

MULTI-SYMPTOM RELIEF

NEW

HALEON

THERAFLU-D

**FLU RELIEF
MAX STRENGTH***

**+ NASAL
DECONGESTANT**

NIGHTTIME

Acetaminophen

Pain Reliever/Fever Reducer

Chlorpheniramine Maleate

Antihistamine

Dextromethorphan HBr

Cough Suppressant

Pseudoephedrine HCl

Nasal Decongestant

**Fast, Powerful,
Multi-Symptom relief of:**

- / Fever
- / Nasal Congestion
- / Sinus Pressure
- / Head & Body Ache
- / Sore Throat Pain
- / Cough
- / Runny Nose
- / Sneezing

**8.0 FL OZ
(237 mL)**

**Cherry
Flavor**

**FAST, POWERFUL
MULTI-SYMPTOM**
relief of:



Fever



Nasal Congestion



Sinus Pressure



Head & Body Ache



Sore Throat Pain



Cough



Runny Nose



Sneezing

*Maximum Strength per 6 hour dose