

Theraflu Flu Relief Maximum Strength Daytime Caplets (cont'd)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- adults and children 12 years of age and over: take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

Other information

- store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients

crospovidone, D&C yellow no. 10 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-855-328-5259

Theraflu Flu Relief Maximum Strength Nighttime Caplets (cont'd)

Drug Facts (continued)

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- adults and children 12 years of age and over: take 2 caplets every 6 hours, while symptoms persist
- do not take more than 6 caplets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	2 caplets every 6 hours
children under 12 years of age	do not use

Other information

- store at controlled room temperature 20-25°C (68-77°F)

Theraflu Flu Relief Maximum Strength Nighttime Caplets (cont'd)

Drug Facts (continued)

Inactive ingredients

crospovidone, FD&C blue no. 2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-855-328-5259

B-0630-770771-10  
REV C



40 TOTAL CAPLETS  
20 DAYTIME CAPLETS  
20 NIGHTTIME CAPLETS

Powerful fever fighting  
Formula that relieves:  
/ Body ache / Headache  
/ Sore throat pain / Cough  
/ Runny nose (Nighttime only)

FLU RELIEF  
MAX STRENGTH\*

Theraflu

NIGHTTIME  
Acetaminophen  
Pain Reliever / Fever Reducer  
Chlorpheniramine Maleate  
Antihistamine  
Dextromethorphan HBr  
Cough Suppressant  
Actual Size  
NT-1

DAYTIME  
Acetaminophen  
Pain Reliever / Fever Reducer  
Dextromethorphan HBr  
Cough Suppressant  
Actual Size  
DT-1  
HALEON

1-855-328-5259

Distributed by:

Haleon

Warren, NJ 07059

Trademarks are owned by or licensed to the Haleon group of companies.

©2022 Haleon group of companies or its licensor.

202019

B-0630-770771-10 REV C

202019

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.  
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.  
DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 6 HOURS AFTER  
THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.  
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.  
READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.

PARENTS:  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org  
TAMPER EVIDENT FEATURE: THERAFLU  
FLU RELIEF MAX STRENGTH CAPLETS ARE  
SEALED IN BLISTER PACKETS. USE ONLY IF  
THE INDIVIDUAL SEAL IS UNBROKEN.

\*Maximum Strength per 6 hour dose.

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.  
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

USE AS DIRECTED

Theraflu Flu Relief Maximum Strength Daytime Caplets

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 500 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
- temporarily reduces fever

Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Theraflu Flu Relief Maximum Strength Nighttime Caplets

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 500 mg	Pain reliever/Fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 15 mg	Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
  - runny nose
- temporarily reduces fever

Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Lift For Additional Drug Facts

Theraflu Flu Relief Maximum Strength Daytime Caplets (cont'd)

Drug Facts (continued)

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product do not use more than directed.

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Theraflu Flu Relief Maximum Strength Nighttime Caplets (cont'd)

Drug Facts (continued)

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children