



Theraflu Flu Relief Max Strength* Daytime Powder (continued)

Drug Facts (continued)

if pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.

Drug Facts (continued)

- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 10 mg, sodium 20 mg
- **phenylketonurics:** contains phenylalanine 19.6 mg per packet
- store at a controlled room temperature at 20 – 25°C (68 – 77°F)

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate

Questions or comments?

1-855-328-5259

Theraflu Flu Relief Max Strength* Nighttime Powder

Drug Facts

Active ingredients (in each packet) **Purposes**
Acetaminophen 1000 mg.....Pain reliever/Fever reducer
Chlorpheniramine maleate 4 mg.....Antihistamine
Dextromethorphan HBr 30 mg.....Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
 - runny nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

- Do not use**
- in a child under 12 years of age
 - if you are allergic to acetaminophen
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains

Drug Facts (continued)

acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days

Theraflu Flu Relief Max Strength* Nighttime Powder (continued)

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**

- take every 6 hours, while symptoms persist. Do not take more than 3 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

Other information

- **each packet contains:** potassium 5 mg, sodium 22 mg
- **phenylketonurics:** contains phenylalanine 12.9 mg per packet
- store at a controlled room temperature at 20 – 25°C (68 – 77°F)

Inactive ingredients acesulfame potassium, anhydrous citric acid, aspartame, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, maltodextrin, natural and artificial flavors, silicon dioxide, sodium citrate, soy lecithin, sucrose, tribasic calcium phosphate

Questions or comments?

1-855-328-5259

**DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME
OR TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.**

PARENTS:
Learn about teen medicine abuse

www.StopMedicineAbuse.org

***Maximum Strength
per 6 hour dose**

**TAMPER EVIDENT INNER UNIT.
DO NOT USE IF SEALED THERAFLU PACKET IS TORN OR BROKEN.**

1-855-328-5259

Distributed by: Haleon, Warren, NJ 07059
Trademarks are owned by or licensed to the Haleon group of companies.
©2022 Haleon group of companies or its licensor. 201926

FLU RELIEF MAX STRENGTH*

M U L T I - S Y M P T O M F L U R E L I E F

HALEON

THERAFLU

FLU RELIEF MAX STRENGTH*

6 x DAYTIME

Acetaminophen Pain Reliever / Fever Reducer
Dextromethorphan HBr Cough Suppressant

6 x NIGHTTIME

Acetaminophen Pain Reliever / Fever Reducer
Chlorpheniramine Maleate Antihistamine
Dextromethorphan HBr Cough Suppressant

Hot liquid therapy that relieves:

- / Fever / Body ache / Headache
- / Sore throat pain / Cough
- / Runny nose (Nighttime only)

**6 DAYTIME PACKETS
6 NIGHTTIME PACKETS
12 TOTAL PACKETS**

**Honey Lemon
Flavor**

USE ONLY AS DIRECTED

**READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.**

**DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.**

**DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN
4 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT
UNLESS DIRECTED BY YOUR DOCTOR.**

Theraflu Flu Relief Max Strength* Daytime Powder

Drug Facts

Active ingredients (in each packet) **Purposes**
Acetaminophen 1000 mg.....Pain reliever/Fever reducer
Dextromethorphan HBr 30 mg.....Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.