

HALEON

# TheraFlu

**FLU RELIEF  
MAX STRENGTH\***

**NIGHTTIME**

**Acetaminophen**

Pain Reliever/Fever Reducer

**Chlorpheniramine Maleate**

Antihistamine

**Dextromethorphan HBr**

Cough Suppressant



**Powerful *fever fighting*  
formula that relieves:**

/ Body ache / Headache  
/ Sore throat pain  
/ Cough / Runny nose

**Honey &  
Elderberry Flavor**

8.3 FL OZ (245.5 mL)



62000000214248

FPO

Drug Facts (continued)

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Drug Facts (continued)

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions • do not use more than directed

- measure the dose correctly using the enclosed dosing cup
- adults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persist
- do not take more than 3 doses (90 mL) in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and over	30 mL
children under 12 years of age	do not use

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.

\*Maximum Strength per 6 hour dose.



Distributed by: Haleon, Warren, NJ 07059 Made in Canada Trademarks are owned by or licensed to the Haleon group of companies. ©2024 Haleon group of companies or its licensor. Pat. Info www.productpats.com

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg ..... Pain reliever/Fever reducer  
Chlorpheniramine maleate 4 mg.....Antihistamine  
Dextromethorphan HBr 30 mg.....Cough suppressant

Purposes

Uses

- temporarily relieves these symptoms due to a common cold or flu:
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation
  - minor sore throat pain
  - runny nose
- temporarily reduces fever

Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

Drug Facts (continued)

If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

**Ask a doctor or pharmacist before use if you are**

Drug Facts (continued)

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product** • avoid alcoholic drinks

- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

Other information

- each 30 mL contains: potassium 5 mg, sodium 34 mg
- store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, glycerin, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Drug Facts (continued)

Questions or comments? 1-855-328-5259

DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD. USE AS DIRECTED