

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME.
DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

Theraflu Flu Relief Maximum Strength* Daytime Syrup (continued)

Drug Facts (continued) Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none">taking the blood thinning drug warfarin Stop use and ask a doctor if <ul style="list-style-type: none">pain or cough gets worse or lasts more than 7 daysfever gets worse or lasts more than 3 daysredness or swelling is presentcough comes back or occurs with rash or headache that lastsnew symptoms occur These could be signs of a serious condition. If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Drug Facts (continued) <ul style="list-style-type: none">do not take more than 3 doses (90 mL) in 24 hours unless directed by a doctorchildren under 12 years of age: do not use <table><tr><td>Age</td><td>Dose</td></tr><tr><td>adults and children 12 years of age and over</td><td>30 mL</td></tr><tr><td>children under 12 years of age</td><td>do not use</td></tr></table> Other information <ul style="list-style-type: none">each 30 mL contains: potassium 5 mg, sodium 34 mgstore at controlled room temperature 20-25°C (68-77°F) Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, glycerin, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate	Age	Dose	adults and children 12 years of age and over	30 mL	children under 12 years of age	do not use
Age	Dose						
adults and children 12 years of age and over	30 mL						
children under 12 years of age	do not use						
Directions <ul style="list-style-type: none">do not use more than directedmeasure the dose correctly using the enclosed dosing cupadults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persist	Questions or comments? 1-855-328-5259						

Theraflu Flu Relief Maximum Strength* Nighttime Syrup

Drug Facts Active ingredients (in each 30 mL) Acetaminophen 1000 mg.....Pain reliever/Fever reducer Chlorpheniramine maleate 4 mg.....Antihistamine Dextromethorphan HBr 30 mg.....Cough suppressant	Purposes Pain reliever/Fever reducer Antihistamine Cough suppressant						
Uses <ul style="list-style-type: none">temporarily relieves these symptoms due to a common cold or flu:<ul style="list-style-type: none">headacheminor aches and painscough due to minor throat and bronchial irritationminor sore throat painrunny nosetemporarily reduces fever							
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none">more than 4,000 mg of acetaminophen in 24 hourswith other drugs containing acetaminophen3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none">skin reddeningblistersrash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.							
Do not use <ul style="list-style-type: none">in a child under 12 years of ageif you are allergic to acetaminophenwith any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have <ul style="list-style-type: none">liver diseaseglaucomatrouble urinating due to an enlarged prostate glanda breathing problem such as emphysema or chronic bronchitiscough that occurs with too much phlegm (mucus)cough that lasts or is chronic such as occurs with smoking, asthma or emphysema	Drug Facts (continued) Ask a doctor or pharmacist before use if you are <ul style="list-style-type: none">taking the blood thinning drug warfarintaking sedatives or tranquilizers When using this product <ul style="list-style-type: none">avoid alcoholic drinksmarked drowsiness may occuralcohol, sedatives and tranquilizers may increase drowsinessbe careful when driving a motor vehicle or operating machineryexcitability may occur, especially in children Stop use and ask a doctor if <ul style="list-style-type: none">pain or cough gets worse or lasts more than 7 daysfever gets worse or lasts more than 3 daysredness or swelling is presentnew symptoms occurcough comes back or occurs with rash or headache that lasts These could be signs of a serious condition. If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.						
	Directions <ul style="list-style-type: none">do not use more than directedmeasure the dose correctly using the enclosed dosing cupadults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persistdo not take more than 3 doses (90 mL) in 24 hours unless directed by a doctorchildren under 12 years of age: do not use <table><tr><td>Age</td><td>Dose</td></tr><tr><td>adults and children 12 years of age and over</td><td>30 mL</td></tr><tr><td>children under 12 years of age</td><td>do not use</td></tr></table> Other information <ul style="list-style-type: none">each 30 mL contains: potassium 5 mg, sodium 34 mgstore at controlled room temperature 20-25°C (68-77°F) Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, glycerin, natural and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate	Age	Dose	adults and children 12 years of age and over	30 mL	children under 12 years of age	do not use
Age	Dose						
adults and children 12 years of age and over	30 mL						
children under 12 years of age	do not use						
	Questions or comments? 1-855-328-5259						

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE. KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

TAMPER EVIDENT INNER UNIT. DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 6 HOURS AFTER THE LAST DOSE OF THE DAYTIME PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

DO NOT TAKE THE DAYTIME AND NIGHTTIME PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

1-855-328-5259

Distributed by: Haleon, Warren, NJ 07059

Trademarks are owned by or licensed to the Haleon group of companies.

©2024 Haleon group of companies or its licensor.

Pat. Info www.productpats.com

See individual boxes for foreign country of origin.



FLU RELIEF MAX STRENGTH* VALUE PACK

MULTI-SYMPTOM FLU RELIEF

HALEON

VALUE PACK



2 - 8.3 FL OZ (245.5 mL) BOTTLES TOTAL 16.6 FL OZ (491 mL) USE AS DIRECTED

*Maximum Strength per 6 hour dose.

**Temporarily controls cough to help you rest. This is not a sleep aid.

Theraflu Flu Relief Maximum Strength* Daytime Syrup

Drug Facts	
Active ingredients (in each 30 mL)	Purposes
Acetaminophen 1000 mg.....	Pain reliever/ Fever reducer
Dextromethorphan HBr 30 mg....	Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold or flu:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
- temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema