

MULTI-SYMPTOM COLD RELIEF

HALEON

# TheraFlu

**SEVERE  
COLD RELIEF**

**+ CHEST CONGESTION**

**Acetaminophen**  
Pain Reliever/Fever Reducer  
**Dextromethorphan HBr**  
Cough Suppressant  
**Guaifenesin / Expectorant**



**Powerful formula** that relieves:

- / Chest congestion
- / Thins and loosens mucus
- / Head and body ache
- / Fever
- / Cough
- / Sore throat pain

8.3 FL OZ (245.5 mL)

**Berry  
Flavor**

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

62000000211337

**Drug Facts** (continued)

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting

**Drug Facts** (continued)

consult a doctor promptly.

**Do not use**

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- with any other product containing diphenhydramine, even one used on skin
- to sedate a child or to make a child sleepy

**Drug Facts** (continued)

- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts** (continued)

**Directions**

- **do not use more than directed**

- measure the dose correctly using the enclosed dosing cup
- take every 4 hours in dosing cup provided, while symptoms persist
- do not take more than 5 doses (150 mL) in 24 hours unless directed by a doctor

Age	Dose
adults and children 12 years of age and over	30 mL
children under 12 years of age	do not use

**Drug Facts** (continued)

**Other information**

- each 30 mL contains: potassium 30 mg, sodium 13 mg
- store at controlled room temperature 20°-25°C (68°-77°F)

**Inactive ingredients**

acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, glycerin, maltitol solution, natural and artificial flavors, propylene glycol, purified water, sodium benzoate, sodium citrate

**Drug Facts** (continued)

**Questions or comments?** 1-855-328-5259

Distributed by: Haleon, Warren, NJ 07059 Made in Canada Trademarks are owned by or licensed to the Haleon group of companies. ©2024 Haleon group of companies or its licensor, Pat. Info www.productpats.com

DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.



**Drug Facts**

Use only as directed

**Active ingredients (in each 30 mL)**

**Purposes**

Acetaminophen 650 mg ..... Pain reliever/Fever reducer  
Diphenhydramine HCl 25 mg ..... Antihistamine/Cough suppressant

**Uses**

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - headache
  - runny nose
  - sneezing
  - itchy nose or throat
- itchy, watery eyes due to hay fever
- cough due to minor throat and bronchial irritation
- temporarily relieves your cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

**Drug Facts** (continued)

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**Drug Facts** (continued)

**When using this product**

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain or cough gets worse or lasts more than 7 days